

1 **Population genomics of *Bombus terrestris* reveals high but unstructured genetic diversity in a potential**
2 **glacial refugium**

3 Sara E. Silva¹, Sofia G. Seabra¹, Luísa G. Carneiro^{1,2}, Vera L. Nunes¹, Eduardo Marabuto¹, Raquel
4 Mendes¹, Ana S. B. Rodrigues¹, Francisco Pina-Martins¹, Selçuk Yurtsever³, Telma G. Laurentino⁴, Elisabete
5 Figueiredo⁵, Maria T. Rebelo⁶, Octávio S. Paulo¹

6
7 ¹Centre for Ecology, Evolution and Environmental Changes (cE3c), Departamento de Biologia Animal,
8 Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal.

9 ²Department of Ecology, Universidade Federal de Goiás, Goiânia, Brazil.

10 ³Biology Department, Science Faculty, Trakya University, Edirne, Turkey.

11 ⁴Zoological Institute, University of Basel, Vesalgasse 1, 4051 Basel, Switzerland.

12 ⁵Linking Landscape, Environment, Agriculture and Food (LEAF), Instituto Superior de Agronomia,
13 Universidade de Lisboa, Tapada da Ajuda, Lisboa, Portugal.

14 ⁶Centre for Environmental and Marine Research (CESAM), Departamento de Biologia Animal, Faculdade
15 de Ciências, Universidade de Lisboa, Lisboa, Portugal.

16
17 **Running title**

18 Population genomics of an Iberian bumblebee.

19
20
21 **Citation:**

22 Sara E Silva, Sofia G Seabra, Luísa G Carneiro, Vera L Nunes, Eduardo Marabuto, Raquel Mendes, Ana S
23 B Rodrigues, Francisco Pina-Martins, Selçuk Yurtsever, Telma G Laurentino, Elisabete Figueiredo, Maria T
24 Rebelo, Octávio S Paulo, Population genomics of *Bombus terrestris* reveals high but unstructured genetic
25 diversity in a potential glacial refugium, *Biological Journal of the Linnean Society*, Volume 129, Issue 2,
26 February 2020, Pages 259–272, <https://doi.org/10.1093/biolinnean/blz182>

30 **Abstract**

31 Ongoing climate change is expected to cause temperature rise and reduction of precipitation levels in the
32 Mediterranean region, which might cause changes in many species' distribution. These effects negatively
33 impact species' gene pools, decreasing genetic variability and adaptive potential. Here we use mitochondrial
34 DNA and RADseq to analyse population genetic structure and genetic diversity of the bumblebee species
35 *Bombus terrestris* (subspecies *B. terrestris lusitanicus*), in the Iberian Peninsula. While this subspecies shows
36 a panmictic pattern of population structure across Iberia and beyond, we found differentiation between
37 subspecies *lusitanicus* and subspecies *africanus*, likely caused by the existence of barriers to gene flow
38 between Iberia and North Africa. Furthermore, results revealed that the Iberian Peninsula harbours a large
39 fraction of *B. terrestris* intraspecific genetic variation, with the highest number of mitochondrial haplotypes
40 found when compared to any other region in Europe studied so far, suggesting a potential glacial *refugium*
41 role for the Iberian Peninsula. Our findings strengthen the idea that Iberia is a very important source of
42 diversity for the global genetic pool of this species, as rare alleles may play a role in population resilience
43 against human or climate-mediated changes.

44
45 **Keywords**

46 Buff-tailed bumblebee, Iberian Peninsula, mtDNA, phylogeography, RADseq.

47

48

49

50

51

52

53

54

55

56

57

58

59 **Introduction**

60 As a consequence of several human-driven environmental changes (e.g. habitat loss, agricultural
61 intensification, use of pesticides, the introduction of new parasites and climate change), insect pollinators
62 have purportedly declined dramatically in recent decades (reviewed in Potts *et al.*, 2010). Moreover, the
63 massive use of managed and commercial bees for crop pollination and honey production (e.g. *Apis mellifera*,
64 *Bombus terrestris*) facilitate the introduction and spread of diseases and parasites (Goulson, 2010), and
65 interfere with the genetic composition of natural populations through hybridization (Jaffé *et al.*, 2016; Seabra
66 *et al.*, 2019).

67 Bumblebees (*Bombus* spp.) have been particularly affected worldwide by the above mentioned problems.
68 Several species have endured dramatic changes in their distribution or became locally extinct in developed
69 regions, such as central and western Europe and North America (Goulson *et al.*, 2008; Bommarco *et al.*,
70 2012). Although several studies have reported high levels of gene flow among European populations
71 (Woodard *et al.*, 2015; Lecocq *et al.*, 2017), some species have experienced a decline on genetic diversity at
72 a local scale (Woodard *et al.*, 2015), which might increase the risk of inbreeding and hinder population's
73 ability to cope with environmental change (Goulson *et al.*, 2008; Maebe *et al.*, 2015).

74 Populations that have persisted in glacial refugia are expected to show higher levels of intraspecific
75 genetic variation comparing to populations outside these regions, due to long-term population persistence
76 and isolation (Hewitt, 1999). The Iberian Peninsula was one of the largest Mediterranean refugia during the
77 Quaternary (2.6 Mya – present) glaciations, and is also at the southernmost latitude limit (rear edge) for
78 many continental species ranges (Hewitt, 2000; Arias *et al.*, 2006). Therefore, many species in Iberia present
79 geographically structured genetic lineages (Miraldo *et al.*, 2011; Rodrigues *et al.*, 2014; Chávez-Galarza *et*
80 *al.*, 2015). Additionally, the proximity of Iberia to Africa, especially at the Strait of Gibraltar and with
81 episodic bridges between the two continents, enabled occasional dispersal of the more vagile organisms,
82 particularly during periods of glacial southern contraction or of postglacial northern expansion (Pinto *et al.*,
83 2013). However, according to Rasmont *et al.* (2015), the Iberian Peninsula is expected to experience major
84 reductions of bumblebee suitable climatic conditions within the forthcoming decades, alongside other
85 southern European regions. Intensive land-use regimes and degradation of semi-natural areas might be
86 already impacting species richness patterns for many taxa across the peninsula (Martins *et al.*, 2014;
87 Newbold *et al.*, 2015). Moreover, the use of commercial bumblebees for crop pollination, which is a

88 common practice in several parts of the peninsula, may negatively impact native natural populations through
89 pathogen spillover and introgression of maladaptive alleles (Murray *et al.*, 2013; Seabra *et al.*, 2019).

90 The buff-tailed bumblebee *Bombus terrestris* (Linnaeus, 1758) (Hymenoptera, Apidae) presents a wide
91 distribution in the West Palearctic region (Rasmont *et al.*, 2008), and has been deliberately introduced as a
92 crop pollinator into several areas worldwide (Ings *et al.*, 2005; Goulson, 2010). Nine subspecies were
93 described based on morphology, particularly in coat colour variation, with additional differences in
94 behaviour, phenology, physiological traits and resistance to parasites (see Figure 1; Rasmont *et al.*, 2008).
95 Mitochondrial and microsatellite variation studies have shown a clear differentiation of northern African and
96 islander *B. terrestris* populations from European mainland ones, with no differentiation amongst the latter. In
97 fact, mainland populations are largely homogeneous with nearly panmixia patterns (Estoup *et al.*, 1996;
98 Widmer *et al.*, 1998; Lecocq *et al.*, 2013b; 2016; Woodard *et al.*, 2015). Genetic cohesiveness found across
99 broad geographical scales has been attributed to the absence of effective barriers to gene flow and
100 bumblebees' great dispersal capability (Estoup *et al.*, 1996; Woodard *et al.*, 2015; Lecocq *et al.*, 2016). Flight
101 radius of *Bombus terrestris* males, for example, varies between 2.6 and 9.9 km (Kraus *et al.*, 2009). Queen
102 dispersal in *B. terrestris* is unknown but estimates for *Bombus pascuorum* (Scopoli, 1763) and *Bombus*
103 *lapidarius* (Linnaeus, 1758) queens show they are able to disperse by at least three and five km, respectively,
104 during their lifetime (Lepais *et al.*, 2010). In the case of *B. terrestris*, two hypotheses have been suggested to
105 explain the genetic homogeneity in European mainland: (1) a recent population expansion from a single
106 glacial refuge, although low sampling efforts hinder definite conclusions (Lecocq *et al.*, 2016); (2) the
107 erosion of past population structure due to genetic homogenisation linked to translocations of commercial *B.*
108 *terrestris*, but this is unlikely since the same pattern of genetic homogenisation was found in the early 1990s,
109 when bumblebee commercialization was barely a practice (Estoup *et al.*, 1996). Unfortunately, studies on *B.*
110 *terrestris* largely undersampled the Iberian Peninsula (see Estoup *et al.* (1996), Widmer *et al.* (1998),
111 Moreira *et al.* (2015) and Lecocq *et al.* (2016)), despite the importance of this area as a potential glacial
112 refuge for the species and its present location at the south-western edge of *B. terrestris* mainland distribution.

113 To address this gap, we explore the role of the Iberian Peninsula in the differentiation dynamics of *B.*
114 *terrestris*, by assessing if (1) Iberian *B. terrestris lusitanicus* (Krüger, 1956) is panmictic with the rest of the
115 *B. terrestris*' distribution, and (2) if Iberian populations of *B. terrestris* harbour standing genetic variation in
116 order to adapt to the currently changing environment. We will address these questions by determining: (i) the

level of genetic differentiation between *B. t. lusitanicus* and other recognised subspecies of *B. terrestris*; (ii) the population genetic structure within the Iberian Peninsula and (iii) the levels of genetic diversity within *B. t. lusitanicus*. We contrast the use of mitochondrial Cytochrome Oxidase I (COI) marker, which has been commonly used to study inter- and intraspecific relationships in bumblebees (Lecocq *et al.*, 2013a, 2016; Dellicour *et al.*, 2015; Moreira *et al.*, 2015), with a novel genome-wide dataset of restriction site-associated DNA sequencing (RADseq). This method readily provides thousands of SNPs and has proven to be effective in biogeography, adaptation, association and conservation studies, even when individual and population sampling is limited (Lozier, 2014; Woodard *et al.*, 2015; Lozier & Zayed, 2016). With RADseq we obtained the first comprehensive dataset of nuclear loci from *B. t. lusitanicus* and evaluated its population structure at a finer scale resolution. This is the first step to investigate the spatial patterns of population structure and genetic diversity on a bumblebee species within the Iberian Peninsula and to identify the main priorities for future research on the Iberian bumblebees' conservation, evolution and environmental adaptation.

Material and Methods

Sampling and DNA extraction

A total of 198 individuals of *B. t. lusitanicus* were collected from six regions within the Iberian Peninsula, covering most of *B. t. lusitanicus*' Iberian distribution and habitat heterogeneity. To minimize the probability of sampling individuals from the same colony, individuals were caught from locations separated by at least one km, within each region, whenever possible (Figure 1B; Table S1, Supporting Information). Additional samples from other subspecies of *B. terrestris* were collected: one *B. t. terrestris* from Switzerland, from Germany and from France, one *B. t. dalmatinus* from Turkey, one *B. t. audax* from Great Britain and four *B. t. africanus* from Morocco (Figure 1A; Table S1, Supporting Information). We focused on diploid individuals (females, mainly workers) for the genetic analyses, so that we would better capture the genetic variation in the populations. The only exception was one male from France, from where we did not collect females.

For the genetic analyses, *B. t. lusitanicus*' samples were grouped according to the six defined regions, while samples from other subspecies were grouped according to the country where they were collected (Table S1, Supporting Information). Thirteen females from closely related species were collected to serve as outgroup: one individual of *Bombus hortorum* (Linnaeus, 1761), three of *B. lapidarius*, three of *Bombus*

146 *lucorum* (Linnaeus, 1761), four of *B. pascuorum* and two of *Bombus ruderatus* (Fabricius, 1775). Samples
147 were preserved for DNA extraction either in absolute ethanol and stored at -20°C, or dried and preserved at -
148 80°C. Total genomic DNA was extracted from fore and mid legs, the head, and for smaller individuals, a
149 portion of the thorax was also used. DNA was isolated with the DNeasy Blood & Tissue extraction kit
150 (Qiagen), following the manufacturer's standard protocol. To maximize DNA yield, some samples were
151 eluted in lower volume of buffer AE (minimum of 60 µl), and the eluted volume was transferred again to the
152 silica column of the kit for a second elution and incubation times with buffer AE were extended up to 30
153 min.

154

155 **COI amplification, RAD libraries construction and sequencing**

156 For all samples, a fragment from the mitochondrial *Cytochrome Oxidase I* gene (COI) was amplified by
157 Polymerase Chain Reaction (PCR) with the primer set LepF/LepR (Hajibabaei *et al.*, 2006). PCR
158 amplifications were carried out in 20 µL volumes containing approximately 10-45 ng of template DNA, 1x
159 reaction buffer, 1.8 mM of MgCl₂, 1.0 mM dNTPs, 0.04 units of *GoTaq Flexi* DNA polymerase and 0.4 µM
160 of each primer. The thermocycling profile consisted of one cycle of one min at 94°C, five cycles of 30 s at
161 94°C, one min at 45°C, and one min at 72°C, followed by 30 cycles of one min at 94°C, one min and 30 s at
162 50°C, and one min at 72°C, with a final step of five min at 72°C. All PCR products were purified with
163 SureClean (Bioline) purification kit and sequenced in the forward direction on an ABI3730XL by Macrogen
164 Europe. DNA sequences were quality controlled with *Sequencher* version 4.0.5 (Gene Codes Corporation).

165 A subset of 55 individuals, including 37 individuals of *B. t. lusitanicus* from the six defined Iberian
166 regions, five individuals from other *B. terrestris* subspecies and 13 individuals from outgroup species (see
167 Figure 1A and B; Table S1, Supporting Information) was used for RAD sequencing analyses. RADseq
168 libraries for Illumina paired-end sequencing were prepared following the protocol by Etter *et al.* (2011)
169 available at <https://www.wiki.ed.ac.uk/display/RADSequencing/Home>, with some minor modifications as
170 reported in Seabra *et al.* (2019). We used the restriction enzyme *Pst*I-HF (New England Biolabs).
171 Sequencing took place on a Illumina HiSeq 2000/2005 at Edinburgh Genomics, Ashworth Laboratories. The
172 55 individuals were run together with other 53 samples from another study (Seabra *et al.*, 2019), over two
173 lanes.

174

COI data analysis

Mitochondrial COI sequences obtained in this study were deposited in GenBank (Accession numbers MN652675 – MN652877; Table S1a, Supporting Information). We also included 17 sequences from our previous study (Seabra *et al.*, 2019) and we followed the designation of haplotypes in that same study (Table S1a, Supporting Information). In order to extend our *B. terrestris* COI dataset (207 sequences) into the species' geographical range, we downloaded from GenBank five additional sets of COI sequences from previous studies, namely Coppée (2010), Williams *et al.* (2012a, b), Moreira *et al.* (2015) and Schmidt *et al.* (2015) (see Figure 1; see Table S1b, Supporting Information, for GenBank accession numbers). In this way, a total of 233 COI sequences were added to our dataset, totalizing 441 sequences. Sequences related to commercially reared or introduced populations were not considered.

COI dataset of *B. terrestris* was aligned in *MAFFT* version 7.271 (Katoh & Standley, 2013) using default settings. The final alignment was checked for accuracy and sequences were trimmed to the same length (597 bp) to eliminate missing data, using *BioEdit* version 7.2.5 (Hall, 1999). Median-joining haplotype networks were constructed in *PopART* version 1.7 (Bandelt *et al.*, 1999; Leigh & Bryant, 2015), in order to visualise the relationship among *B. terrestris* haplotypes. We used *Arlequin* version 3.5.2.2 (Excoffier & Lischer, 2010) and only the individuals from the Iberian Peninsula to perform a standard analysis of molecular variance (AMOVA) with 10,000 permutation steps and calculate haplotype (h) and nucleotide (π) diversities. We also calculated haplotype and nucleotide diversities for the individuals of the remaining *B. terrestris* distribution range in order to compare with the Iberian Peninsula. File format conversion for *PopART* and *Arlequin* were performed using *PGDSpider* version 2.1.0.3 (Lischer & Excoffier, 2012).

RADseq data analysis

RADseq data obtained in this study are available at Sequence Read Archive (PRJNA578045). All console commands used for RADseq data filtering, SNP discovery and subsequent analyses are available in Appendix 2. Quality control of the RADseq raw read data was performed using *FastQC* version 0.11.3 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). The script *process_radtags* implemented in *Stacks* version 1.29 (Catchen *et al.*, 2013) was used (with default settings) to remove low quality (with a phred quality score below 33) and unidentifiable reads as well as to demultiplex the data. *Bowtie2* version 2.1.0 (Langmead & Salzberg, 2012) was used to trim the last three bases from the 3' end of each read before

alignment, as this region is richer in low-quality base calls, and to align each sample's reads to the assembled reference genome of *B. terrestris* (NCBI Assembly GCA_000214255.1) with the "sensitive" option. *SAMtools* version 0.1.19 (Li *et al.*, 2009) was used to remove low quality alignments (mapping quality below 20) and any unmapped reads.

After data filtering, RAD loci were identified and SNPs called using the *Stacks* pipeline 1.45 (Catchen *et al.*, 2013). In order to minimize the impact of differences among taxa in the number of SNPs obtained and the amount of missing data, three datasets were created from the initial 55 individuals for different analyses (see Table S1, Supporting Information): (1) BT_OUT: includes all the studied subspecies (all *B. terrestris*) as well as the outgroup species; (2) BT_SSP: includes all *B. terrestris* subspecies and does not include the outgroup species; (3) BT_BTL: includes the Iberian *B. t. lusitanicus* only.

Preliminary tests were carried out to identify optimal *Stacks* parameters (Appendix 1, Figures S3, S4, S5, Supporting Information). In *pstacks*, minimum stack depth was set to six; in *cstacks*, maximum number of mismatches allowed when building catalog loci was two; in *populations*, individuals were grouped into putative locations based on geographical region (see Table S1, Supporting Information) and one random SNP per RAD locus was used to avoid confounding signals of linkage disequilibrium. SNPs were only retained if they were present in 50% of individuals in at least n-1 (where n is the number of geographic regions) in all datasets. To test for differences among outputs with different randomly selected SNPs, module *populations* was run several times, but no significant differences were obtained in the final results (data not shown).

Finally, we performed an additional filtering step to remove loci with minor allele frequencies <0.05 and more than 25% of missing data across all samples using *VCFTools* version 0.1.15 (Danecek *et al.*, 2011). *VCFTools* was also used to obtain the mean coverage per site per individual. The resulting SNP datasets were then used in subsequent population genomic and phylogeographic analyses: dataset BT_OUT to check for differentiation between *B. terrestris* and outgroup species; dataset BT_SSP to determine differentiation between *B. t. lusitanicus* and the other subspecies; dataset BT_BTL to determine population genetic structure and genetic diversity of *B. t. lusitanicus* within the Iberian Peninsula.

Principal component analyses (PCA) of the three RADseq datasets were performed using the package SNPRelate version 1.12.0 (Szulkin *et al.*, 2016) as implemented in the R script *snp_pca_static.R* (https://github.com/CoBiG2/RAD_Tools) as of commit "bb2fc45". To test for differentiation between *B. t. lusitanicus* and the other subspecies and within *B. t. lusitanicus*, clustering analyses of population structure

233 were performed using the datasets BT_SSP and BT_BTL and *MavericK* version 1.0.4 (Verity & Nichols,
234 2016). As the model used by *MavericK* assumes that markers are neutral (Verity & Nichols, 2016), we first
235 performed an outlier analysis of both datasets to identify and remove non-neutral markers, using both
236 *BayeScan* version 2.1 (Foll & Gaggiotti, 2008) and *SelEstim* version 1.1.7 (Vitalis *et al.*, 2014). *BayeScan*
237 was run using a matrix of SNP genotypes, with prior odds for the neutral model turned to five and assuming
238 a detection threshold of 0.05. The remaining parameters were set to default values. Plots and convergence
239 were checked using the *R* script *plot_R.r* available within the *BayeScan* package, and the package CODA
240 version 0.19-1 (Plummer *et al.*, 2006). *SelEstim* was run after randomizing the reference allele for each locus
241 (using the *R* script *SelEstim.R* available within *SelEstim* package) and using 50 pilot runs of 1 K length,
242 followed by a main run of 1 M length with a burnin of 100 K, a thinning interval of 20, and a detection
243 threshold of 0.01. The *R* script *SelEstim.R* was also used to obtain the list of outliers and check for
244 convergence. The total number of outliers identified by *BayeScan* and *SelEstim* were removed from the
245 datasets using the *Python* script *outlier_removal.py* (https://github.com/CoBiG2/RAD_Tools) as of commit
246 “ba731f2”. Datasets with only neutral markers were analysed using *MavericK* version 1.0.4 (Verity &
247 Nichols, 2016), wrapped under *Structure_threader* version 1.2.4 (Pina-Martins *et al.*, 2017) for values of “K”
248 between one and five for dataset BT_SSP, and values between one and four for the dataset BT_BTL. We first
249 performed a single “pilot” run of 5,000 iterations, with a burnin of 500 using an admixture model, a free
250 alpha parameter of one with a standard deviation of the normal proposed distribution of 0.10 and
251 “thermodynamic integration” (TI) turned off. Posterior median and posterior standard deviation of alpha
252 were obtained from the “pilot” run and used in a “tuned” run as parameters for the admixture model as
253 follows: alpha was set to 10 times the posterior median and alphaPropSD to five times the posterior SD. This
254 “tuned” run was comprised of five runs of 10,000 iterations, with a burnin of 2000, with TI turned on and
255 another set of 20 runs of 10,000 samples with a burnin of 2000. The most suitable value of “K” was
256 calculated for both datasets using the TI method. The *R* script *MavericK1.0_functions.R* available within the
257 *MavericK* package was used to produce diagnostic plots in order to check for convergence and
258 autocorrelation, and the Qmatrix plots.

259 Finally, a locus-by-locus AMOVA was performed in *Arlequin* with RADseq dataset BT_BTL, which
260 includes all the SNPs, in order to examine the variance within and among geographical regions, and
261 significance was calculated using 10,000 permutation steps. Genome-wide measures of genetic diversity,

including per-SNP nucleotide diversity (π_{SNP}), and the mean per-individual observed and expected heterozygosities (H_o and H_e) were calculated using *VCFTools* and the same dataset.

File format conversions for *BayeScan*, *MaverickK* and *Arlequin* were performed using *PGDSpider* whereas file format conversion for *SelEstim* was performed with the bash script *GESTE2SelEstim.sh* (https://github.com/Telpidus/omics_tools) as of commit “f74f66b”.

Results

The final alignment of the mitochondrial COI dataset included no indels, and consisted of a total of 26 variable sites, of which nine were parsimony-informative. A total of 16 haplotypes were found (Table S1a and S1b, Supporting Information; Figure 2).

After Illumina sequencing of RAD libraries for 55 individuals, we obtained an average of 7.7 M paired end reads of 125 bp, per individual. Of those, an average of 7.4 M were retained after filtering with *process_radtags*, representing approximately 8.18% of the genome sequenced. Quality scores of the retained reads ranged from 36 to 37, with a GC content of ~40%. An average of 52.69% of the quality-filtered reads aligned to the *B. terrestris* genome. Of the 55 sequenced individuals, nine were excluded from further analyses (six *B. t. lusitanicus* from four of the Iberian defined regions and three representatives of the outgroup species) due to lower mean coverage per individual ($\leq 12\times$). Missing data of the remaining samples averaged 5.4%. Information concerning the output of RADseq filtering steps for each sample (number of initial and mapped reads, coverage, datasets at which samples belong and missing data) are provided in Table S1 (Supporting Information). Final individual counts and statistics per dataset after filtering were as follows: BT_OUT dataset comprises 46 individuals and 5,357 SNPs (aprox. 22.7 markers/ Mb) with a mean coverage of 57.7X per site, per individual; BT_SSP dataset comprises 36 individuals and 10,765 SNPs (aprox. 45.6 markers/ Mb) with a mean coverage of 52.3X per site, per individual; BT_BTL dataset comprises 31 individuals and 11,369 SNPs (aprox. 48.2 markers/ Mb) with a mean coverage of 56.6X per site, per individual.

Differentiation of *Bombus terrestris lusitanicus*

Seven of the 16 *B. t. lusitanicus*’ COI haplotypes are present in the Iberian Peninsula. The most common haplotype (H1) is common to the entire Peninsula and to the remaining European mainland regions analysed

291 whereas H2 is shared with central Europe (Switzerland) and Ireland (Figure 2). The other five haplotypes
292 found in Iberia are exclusive to this peninsula (Figure 2). No shared haplotypes were detected between the
293 Iberian Peninsula and North Africa. Haplotypes H9 and H16 are exclusive of North Africa and H14 is shared
294 between this region and Italy. Some haplotypes found in Great Britain, Sardinia, Corsica and Canary Islands
295 are also exclusive from these regions (Figure 2A). Most of the haplotypes differ from H1 in only one or two
296 mutational steps, with the exception of the haplotypes found only in islands or in North Africa, the
297 haplotypes from Sardinia and Morocco being the most differentiated (Figure 2C).

298 PCA of RADseq dataset BT_OUT revealed a clear separation between *B. terrestris* and the outgroup
299 species, with the exception of *B. lucorum* which is the closest to *B. terrestris* (Figure 3A). PCA using dataset
300 BT_SSP, with *B. terrestris* samples only, revealed a very narrow separation between samples from the
301 Iberian Peninsula and those from Great Britain and Germany across EV2 (EV2 explains 3.08% of the
302 variance). On the other hand, individuals from Morocco show a greater separation across EV1 from the
303 remaining samples (EV1 explains 10.14% of the variance; Figure 3B). Outlier tests of dataset BT_SSP
304 revealed that a total of 44 SNPs were non-neutral: 43 SNPs (0.40% of the total SNPs) when using *SelEstim*,
305 and two SNPs (0.02% of the total SNPs) when using *BayeScan* (one SNP was identified by both softwares).
306 Clustering analyses of the dataset BT_SSP with non-neutral loci removed, using *MaverickK*, determined the
307 existence of two groups ($K = 2$) as the most likely scenario (Figure 3C). These two groups correspond to i)
308 the individuals from the Iberian Peninsula, Great Britain and Germany, and ii) individuals from Morocco,
309 which is consistent with what was observed in the PCA (Figure 3B).

310

311 **Iberian populations genetic structure**

312 The most common COI haplotype H1 is present in 95% of the *B. terrestris* sampled in the Iberian
313 Peninsula. H2 is present in three samples from IP-NO and IP-WE regions, while H3 is only present in two
314 samples from southern locations, one IP-SE and another from IP-SW, respectively. All the remaining Iberian
315 COI haplotypes (H4-H7) are represented by a single sample (Figure 2B). The Iberian region with the highest
316 haplotype diversity is IP-SE with five haplotypes (Figure 2B and Table S2, Supporting Information). The
317 haplotype network did not show a structured phylogeographic pattern, with the most common haplotype
318 being shared among geographically distant regions, and the less frequent and unique haplotypes being
319 closely related to H1, in a 'star-like' configuration (Figure 2C).

PCA of RADseq dataset BT_BTL showed no clear separation between any Iberian geographic regions (EV1 and EV2 only explain 3.75% and 3.67% of the variation, respectively; Figure 3D). The most segregated individuals are BTL_075 (IP-NW) and BTL_136 (IP-SE) along EV1, and BTL_306 (IP-WE) along EV2. Missing data values for these individuals does not explain their separation from the remaining (3.7%, 5.7% and 10.6% respectively), and their COI haplotype is the most common one, H1. Outlier tests of dataset BT_BTL revealed a total of 31 non neutral SNPs: 29 (0.26% of the total SNPs) when using *SelEstim*, and two SNPs (0.02% of the total SNPs) when using *BayeScan*. Clustering analyses of the dataset BT_BTL with non-neutral loci removed, using *MaverickK*, determined the existence of one group ($K = 1$) as the most likely scenario (data not shown), which is concordant with the PCA results.

Genetic diversity

AMOVA results revealed an absence of genetic structure for *B. terrestris* COI dataset, suggesting that the overall source of variation was within geographical regions instead of among these (Table 1). Measures for haplotype (h) and nucleotide (π) diversities calculated using *B. terrestris* COI dataset for the total Iberian Peninsula were 0.08470 and 0.00018, respectively (Table 2). More than 43% (7 in 16) of the haplotypes found were present in the Iberian Peninsula, and 31% (5 in 16) were exclusive of this area and no other European region showed such a high number of haplotypes. It is important to note, however, that sample sizes differ substantially (Figure 2 and Table 2).

AMOVA results using the RADseq dataset BT_BTL mirror those of the COI dataset, with an absence of genetic structure, indicating that the overall source of variation is within and not among geographical regions (Table 1). Measures of per-SNP nucleotide diversity and mean per-individual observed and expected heterozygosities for the total Iberian Peninsula were 0.2780, 0.2326 and 0.2773, respectively (Table 2).

Discussion

We conducted a population genetic study, with mitochondrial and nuclear genome-wide markers, to measure differentiation of *B. t. lusitanicus* from other *B. terrestris*' subspecies, and to investigate this subspecies' population structure and genetic diversity within the Iberian Peninsula. We found no evident differentiation pattern on mitochondrial DNA between *B. t. lusitanicus* and the other European mainland subspecies *B. t. terrestris*, *B. t. dalmatinus* and *B. t. calabricus*. On the other hand, we found a clear

349 differentiation of North African *B. t. africanus* from the remaining subspecies, including the geographically
350 close *B. t. lusitanicus*, which is in accordance with previous studies (Coppée, 2010; Lecocq *et al.*, 2016) and
351 confirmed here with samples from southern Iberia, where any evidence of admixture with *B. t. africanus*
352 would be more likely to occur. Considering the genetic diversity of *B. terrestris* at the species level, it is
353 homogeneous across mainland populations, while subspecies from the islands and North Africa appear to be
354 more differentiated, although with evidence of some admixture, particularly between the British and
355 continental populations (as previously reported by Moreira *et al.*, 2015). The presence of the same haplotype
356 in both Algeria and central Italy could also indicate some admixture in this region. Our study corroborates
357 those of Estoup *et al.* (1996), Widmer *et al.* (1998), Moreira *et al.* (2015) and Lecocq *et al.* (2016) based on
358 mitochondrial and microsatellite markers. Our genome-wide analyses with RAD sequencing show very
359 slight distinction of *B. t. lusitanicus* from *B. t. terrestris* (Germany) and *B. t. audax* (Great Britain) samples
360 in the PCA, although not supported by *Maverick* results, but corroborates the clear differentiation from *B. t.*
361 *africanus*. However, larger sample sizes of *B. t. terrestris* and *B. t. audax* are needed to better evaluate this
362 small differentiation at the genomic level.

363 The lack of differentiation of *B. t. lusitanicus* from other European mainland subspecies can be explained
364 by: (1) a common origin with subsequent local differentiation, which is supported by the star-like pattern in
365 the mitochondrial COI network with rarer haplotypes deriving from a single ancestral haplotype (H1); (2)
366 high dispersal ability of these insects across large distances (Kraus *et al.*, 2009; Lepais *et al.*, 2010) and
367 extensive mountain ranges such as the Pyrenees; or (3) erosion of genetic differentiation caused by
368 hybridization with commercial hives from allochthonous origin which are used in several areas in Europe for
369 crop pollination (commercial stocks used in the Iberian Peninsula include mostly subspecies *B. t. terrestris*
370 and *B. t. dalmatinus* (Lecocq *et al.*, 2016; Velthuis & van Doorn, 2006)). Putative hybrids with commercial
371 hives in the western Iberian Peninsula were already detected (Seabra *et al.*, 2019), but a widespread genetic
372 erosion is not expected because the transfer of colonies of this species across Europe for crop pollination is a
373 relatively recent phenomena (Estoup, 1996).

374 On the other hand, the differentiation found between *B. t. lusitanicus* and *B. t. africanus* suggests that
375 gene flow between the Iberian Peninsula and North Africa is much lower than to elsewhere in mainland
376 Europe. The number of accumulated differences on mitochondrial DNA also suggests these two subspecies
377 probably started to diverge earlier than the others. The Mediterranean sea thus seem to be an effective barrier

378 to gene flow, even though the two continents are geographically very close at the Strait of Gibraltar (< 15 km
379 at the closest point) and despite the good dispersal capability of bumblebees. According to the information
380 retrieved from <http://www.atlashymenoptera.net/>, *B. terrestris* is currently present right up to the coast on
381 both sides of the Strait. Both Moreira *et al.* (2015) and Estoup *et al.* (1996) reported evidence of *B. terrestris*
382 migrating over sea, across the English Channel and between the Isle of Man and Ireland, though at
383 recognisably very low rates. Also other bumblebee species, such as *Bombus jonellus* (Kirby, 1802) was
384 found to be able to disperse over sea barriers up to 30 km (Darvill *et al.*, 2010). As bumblebees are known to
385 disperse such large distances, two hypothesis might explain this result: 1) migration is conditioned by the
386 prevailing wind conditions, characterised by strong winds from easterly or westerly directions, which are
387 known to have an important role in dispersion patterns of several species (e.g. moth *Cornifrons ulceratalis*
388 (Dantart *et al.*, 2009); 2) migration occurs between both continents, but local differences in environmental
389 conditions, and/or sexual selection, may be acting against migrants and preventing effective gene flow. The
390 Strait of Gibraltar seems to hinder the dispersal of other flying species between North Africa and Iberian
391 Peninsula (e.g. the butterfly *Pararge aegeria* (Weingartner *et al.*, 2006)) while acted as a route of dispersal
392 for others, mainly during lower sea level periods (e.g. the Iberian honey bee *Apis mellifera iberiensis*
393 (Chávez-Galarza *et al.*, 2015)).

394 We did not find population genetic structure within *B. t. lusitanicus* across the Iberian Peninsula,
395 contrarily to what was reported for the Iberian honey bee *A. m. iberiensis* (Chávez-Galarza *et al.*, 2015). In
396 this latter species, two highly divergent genetic lineages are observed which form a northeastern-
397 southeastern cline, better explained by secondary contacts between divergent populations from distinct and
398 isolated glacial refugia (Chávez-Galarza *et al.*, 2015). The panmictic pattern of *B. t. lusitanicus* within Iberia
399 is likely due to *B. terrestris*' long-distance flights capability, coupled with the absence of effective
400 geographical barriers to its dispersal. Long-distance flights of queens and males contribute towards regular
401 gene flow and were suggested to be sufficient to maintain genetic cohesion of common bumblebees' species
402 over large areas (Lepais *et al.*, 2010). Also, the fact that *B. terrestris* is a short-tongued generalist bumblebee
403 (Chapman *et al.*, 2003), having a large foraging range (Walther-Hellwig & Frankl, 2000), probably increases
404 its capacity to find suitable habitats under a variety of conditions. In addition, this species is tolerant to a
405 broad range of climates (Penado *et al.*, 2016), from Mediterranean beaches under high temperatures to high
406 mountains of crio-oromediterranean regimes. Thus, individuals are more capable to disperse and occupy a

407 vast majority of habitats when compared to other species, contributing to the observed large-scale
408 connectivity. The already referred hypothesis of a recent population expansion from a single periglacial
409 refuge in the Iberian Peninsula could also explain not only the observed pattern of homogeneity but also the
410 star-like pattern in the mitochondrial COI network. This refuge may have existed in the Iberian Peninsula or
411 elsewhere in Europe (as also referred by Estoup *et al.*, 1996), as the most common haplotype H1 is
412 widespread across the continent. Nonetheless, some rarer haplotypes could have evolved in a smaller refuge
413 in the Betic ranges of southern Spain ('refugia within refugia' paradigm of Gómez & Lunt (2007)), which
414 might explain the higher genetic diversity found in mitochondrial DNA for the region IP-SE (Figure 2B;
415 Table S2). This region is characterized by semi-arid low-lands drastically contrasting with steep changes in
416 vegetation and climate along an altitudinal cline. It is considered as a hotspot for Mediterranean biodiversity,
417 harbouring many endemic species or lineages, and so the high genetic diversity found in this region (five
418 haplotypes from a total of 10 in continental Europe) is not unexpected (Hewitt, 2011; Nunes *et al.*, 2014).
419 The hypothesis of admixture between this region and North Africa is unlikely because there are no
420 haplotypes shared between both regions as referred before. The use of commercial hives of *B. terrestris* for
421 crop pollination in Southeast of Spain (IP- SE), and in IP-SW and IP-WE, is a common practice (Cejas *et al.*,
422 2018). In these regions, commercial bumblebees have been found foraging outside greenhouses and on
423 natural habitats, and introgression between commercial and native bumblebees was detected (Cejas *et al.*,
424 2018, 2019; Seabra *et al.*, 2019; Trillo *et al.*, 2019). Thus some of the variation found in IP-SE could have
425 been artificially introduced. We found one COI haplotype (H3) in regions IP-SE and IP-SW which is also
426 relatively common in commercial stocks and in individuals collected from greenhouse areas investigated by
427 Seabra *et al.* (2019). This haplotype was found in two specimens: one collected from about 300 km from the
428 area where Cejas *et al.* (2018, 2019) detected potential hybrids between commercial and native populations,
429 based on morphological and mitochondrial 16S data; one collected from about 30 km from the area where
430 Seabra *et al.* (2019) also detected potential introgression between both and escaped individuals, based on
431 RADseq data. None of the other unique haplotypes from IP_SE were found within the commercial samples
432 analysed by Seabra *et al.* (2019). Moreira *et al.* (2015) found that commercially reared populations were
433 differentiated from the majority of the wild populations from Ireland, having the highest number of private
434 microsatellite alleles. Thus the introduction of non-native specimens can lead to changes in the genetic
435 structure of the native ones, and ultimately, increase the risk of displacement and the consequent loss of rare

436 beneficial alleles, especially in populations with low genetic diversity. In this way, we cannot underestimate
437 the potential impact of the use of allochthonous commercial bumblebees for local crop pollination,
438 independent of its origin.

439 The extended sampling in the Iberian Peninsula revealed this region as one of the richest in genetic
440 diversity for *B. terrestris*, with the highest number of mitochondrial COI haplotypes than any other region in
441 Europe studied so far (though some mainland regions remain undersampled). Iberia seems to be an important
442 source of diversity for the global genetic pool of this species, as rare alleles may play a role in population
443 resilience against human or climate-mediated changes (Barret & Schluter, 2007), especially at the extremes
444 of the species range. Since this is the first study evaluating population genetic diversity in *B. terrestris* with
445 RADseq, we were not able to compare our results with other regions in Europe regarding the diversity at the
446 genomic level. When comparing with North American *Bombus* species (Lozier, 2014; Jackson *et al.*, 2018),
447 despite the different RADseq markers used, *B. t. lusitanicus* from the Iberian Peninsula showed similar or
448 higher diversity values, even when comparing with *Bombus impatiens* Cresson, 1863, a common species in
449 eastern North America.

450 Further ecological studies comparing habitats, phenology and phenotypic characteristics of *B. terrestris*
451 from the south of the peninsula with those from North Africa could help to describe barriers to dispersion
452 and to gene flow within this species in more detail. Also, the absence of population genetic structure will
453 facilitate the analyses of the adaptive potential of *B. t. lusitanicus* to environmental changes within the
454 Iberian Peninsula, by finding adaptive genetic diversity and by modelling species response to future land-use
455 and/or climatic changes. Ecological and genetic studies focusing in arid regions or in other Mediterranean
456 peninsulas are also needed, in order to understand how environmental change is affecting natural populations
457 of *B. terrestris*. It could also help to evaluate impacts of global warming on crop pollination efficiency of
458 commercial hives at the extremes of this species range and in economically important regions.

459

460 **Acknowledgements**

461 This work was funded by national funds through FCT – Fundação para a Ciência e a Tecnologia (project
462 UID/BIA/00329/2013 (2015-2018); UID/AGR/04129/2013; grant PD/BD/113548/2015, under the PhD
463 program “Biology and Ecology of Global Changes”, Univ. Aveiro & Univ. Lisbon, Portugal, attributed to
464 S.E.S.; grant SFRH/BPD/108413/2015, attributed to S.G.S.) and through ProDer – Programa de

Desenvolvimento Rural (Project ref. ProDeR 4.1. 46221-3). Thanks are due for the financial support also to CESAM (UID/AMB/50017/2019), to FCT/MCTES through national funds, and the co-funding by the FEDER, within the PT2020 Partnership Agreement and Compete 2020. We thank Edinburgh Genomics, particularly Karim Gharbi, for their assistance with RAD sequencing and Moises Mallo (IGC) for the use of Bioruptor. We also thank the anonymous referees for helpful comments on an earlier version of the manuscript.

471

472 **Author's contributions**

SES, SGS, LGC and OSP designed the study. SES, VLN, RM, ASB, EM, SY, TGL, EF, JM, MTR and OSP were responsible for sampling. SES, SGS and VLN were responsible for DNA extraction and mitochondrial DNA amplification. VLN constructed RAD libraries. SES and SGS performed the bioinformatic analyses, with important contributions from VLN, FPM and OSP. SES wrote the manuscript with contributions from all the other authors.

478

479 **References**

- 480 Arias MC, Rinderer TE, Sheppard WS. 2006. Further characterization of honey bees from the Iberian
481 Peninsula by allozyme, morphometric and mtDNA haplotype analyses. *Journal of Apicultural Research*
482 45: 188–196.
- 483 Bandelt H-J, Forster P, Rohl A. 1999. Median-Joining networks for inferring intraspecific phylogenies.
484 *Molecular Biology* 16: 37–48.
- 485 Barret RDH, Schluter D. 2008. Adaptation from standing genetic variation. *Trends in Ecology and Evolution*
486 23: 38-44.
- 487 Bommarco R, Lundin O, Smith HG, Rundlof M. 2012. Drastic historic shifts in bumble-bee community
488 composition in Sweden. *Proceedings of the Royal Society B: Biological Sciences* 279: 309–315.
- 489 Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. 2013. Stacks: An analysis tool set for
490 population genomics. *Molecular Ecology* 22: 3124–3140.

491 Cejas D, López-López A, Muñoz I, Ornos C, De la Rua P. 2019. Unveiling introgression in bumblebee
492 (*Bombus terrestris*) populations through mitogenome-based markers. *Animal Genetics*.

493 Cejas D, Ornos C, Muñoz I, De la Rua P. 2018. Searching for molecular markers to differentiate *Bombus*
494 *terrestris* (Linnaeus) subspecies in the Iberian Peninsula. *Sociobiology* 65: 558-565.

495 Chapman RE, Wang J, Bourke AFG. 2003. Genetic analysis of spatial foraging patterns and resource sharing
496 in bumble bee pollinators. *Molecular Ecology* 12: 2801–2808.

497 Chávez-Galarza J, Henriques D, Johnston JS, Carneiro M, Rufino J, Patton JC, Pinto MA. 2015. Revisiting
498 the Iberian honey bee (*Apis mellifera iberiensis*) contact zone: Maternal and genome-wide nuclear
499 variations provide support for secondary contact from historical refugia. *Molecular Ecology* 24: 2973–
500 2992.

501 Coppée A. 2010. *Bombus terrestris* (L. 1758): A complex species or a species complex? -
502 Intraspecific pheromonal and genetic variations of *Bombus terrestris* (L.). Impacts on the
503 speciation. PhD thesis. Université de Mons, Laboratoire de Zoologie.

504 Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT,
505 Sherry ST, McVean G, Durbin R, 1000 Genomes Project Analysis Group. 2011. The variant call format
506 and VCFtools. *Bioinformatics* 27: 2156–2158.

507 Dantart J, Stefanescu C, Àvila A, Alarcón M. 2009. Long-distance wind-borne dispersal of the moth
508 *Cornifrons ulceratalis* (Lepidoptera: Crambidae: Evergestinae) into the northern Mediterranean.
509 *European Journal of Entomology* 106: 225-229.

510 Darvill B, O'Connor S, Lye GC, Waters J, Lepais O, Goulson D. 2010. Cryptic differences in dispersal lead
511 to differential sensitivity to habitat fragmentation in two bumblebee species. *Molecular Ecology* 19:
512 53–63.

513 Dellicour S, Michez D, Mardulyn P. 2015. Comparative phylogeography of five bumblebees: impact of
514 range fragmentation, range size and diet specialization. *Biological Journal of the Linnean Society* 116:
515 926–939.

516 Estoup A, Solignac M, Cornuet JM, Goudet J, Scholl A. 1996. Genetic differentiation of continental and
517 island populations of *Bombus terrestris* (Hymenoptera: Apidae) in Europe. *Molecular Ecology* 5: 19–
518 31.

519 Etter PD, Bassham S, Hohenlohe PA, Johnson EA, Cresko WA. 2011. SNP Discovery and genotyping for
520 evolutionary genetics using RAD sequencing. *Methods in Molecular Biology* 772: 157–178.

521 Excoffier L, Lischer H. 2010. Arlequin suite ver 3.5: A new series of programs to perform population
522 genetics analyses under Linux and Windows. *Molecular Ecology Resources* 10: 564–567.

523 Foll M, Gaggiotti O. 2008. A genome-scan method to identify selected loci appropriate for both dominant
524 and codominant markers: A Bayesian perspective. *Genetics* 180: 977–993.

525 Gómez A, Lunt DH. 2007. Refugia within refugia: patterns of phylogeographic concordance in the
526 Iberian Peninsula. In: Weiss S, Ferrand N, eds. *Phylogeography of southern European refugia*.
527 The Netherlands: Springer, 155–188.

528 Goulson D. 2010. Impacts of non-native bumblebees in Western Europe and North America. *Applied*
529 *Entomology and Zoology* 45: 7–12.

530 Goulson D, Lye GC, Darvill B. 2008. Decline and conservation of bumble bees. *Annual Review of*
531 *Entomology* 53: 191–208.

532 Hajibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PDN. 2006. DNA barcodes distinguish species
533 of tropical Lepidoptera. *Proceedings of the National Academy of Sciences* 103: 968–971.

534 Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for
535 Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.

536 Hewitt GM. 1999. Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society*
537 68: 87–112.

538 Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405: 907–913.

539 Hewitt GM. 2011. Mediterranean peninsulas: the evolution of hotspots. In: Zachos FE, Habel JC, eds.
540 *Biodiversity hotspots*. Amsterdam, The Netherlands: Springer, 123–147.

541 Ings TC, Schikora J, Chittka L. 2005. Bumblebees, humble pollinators or assiduous invaders? A population
542 comparison of foraging performance in *Bombus terrestris*. *Oecologia* 144: 508–516.

543 Jackson JM, Pimsler ML, Oyen KJ, Koch-Uhuad JB, Herndon JD, Strange JP, Dillon ME, Lozier JD. 2018.
544 Distance, elevation and environment as drivers of diversity and divergence in bumble bees across
545 latitude and altitude. *Molecular Ecology* 27: 2926–2942.

546 Jaffé R, Pope N, Acosta AL, Alves DA, Arias MC, De la Rúa P, Francisco FO, Giannini TC, González-
547 Chaves A, Imperatriz-Fonseca VL, Tavares MG, Jha S, Carnevalheiro LG. 2016. Beekeeping practices
548 and geographic distance, not land use, drive gene flow across tropical bees. *Molecular Ecology* 25:
549 5345–5358.

550 Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in
551 performance and usability. *Molecular Biology and Evolution* 30: 772–780.

552 Kraus FB, Wolf S, Moritz RFA. 2009. Male flight distance and population substructure in the bumblebee
553 *Bombus terrestris*. *Journal of Animal Ecology* 78: 247–252.

554 Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 9: 357–359.

555 Lecocq T, Coppée A, Michez D, Brasero N, Rasplus J-Y, Valterová I, Rasmont P. 2016. The alien's identity:
556 consequences of taxonomic status for the international bumblebee trade regulations. *Biological*
557 *Conservation* 195: 169–176.

558 Lecocq T, Dellicour S, Michez D, Lhomme P, Vanderplanck M, Valterová I, Rasplus J-Y, Rasmont P. 2013a.
559 Scent of a break-up: phylogeography and reproductive trait divergences in the red-tailed bumblebee
560 (*Bombus lapidarius*). *BMC Evolutionary Biology* 13: 263.

561 Lecocq T, Gérard M, Michez D, Dellicour S. 2017. Conservation genetics of European bees: new insights
562 from the continental scale. *Conservation Genetics* 18: 585–596.

563 Lecocq T, Vereecken NJ, Michez D, Dellicour S, Lhomme P, Valterová I, Rasplus J-Y, Rasmont P. 2013b.
564 Patterns of genetic and reproductive traits differentiation in mainland vs. Corsican populations of
565 bumblebees. *PLoS ONE* 8: e65642.

566 Leigh JW, Bryant D. 2015. PopART: Full-feature software for haplotype network construction. *Methods in*
567 *Ecology and Evolution* 6: 1110–1116.

568 Lepais O, Darvill B, O'Connor S, Osborne JL, Sanderson RA, Cussans J, Goffe L, Goulson D. 2010.
569 Estimation of bumblebee queen dispersal distances using sibship reconstruction method. *Molecular*
570 *Ecology* 19: 819–831.

571 Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome
572 Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools.
573 *Bioinformatics* 25: 2078–2079.

574 Lischer HEL, Excoffier L. 2012. PGDSpider: An automated data conversion tool for connecting population
575 genetics and genomics programs. *Bioinformatics* 28: 298–299.

576 Lozier JD. 2014. Revisiting comparisons of genetic diversity in stable and declining species: assessing
577 genome-wide polymorphism in North American bumble bees using RAD sequencing. *Molecular*
578 *Ecology* 23: 788–801.

579 Lozier JD, Zayed A. 2017. Bee conservation in the age of genomics. *Conservation Genetics* 18: 713–729.

580 Maebe K, Meeus I, Ganne M, De Meulemeester T, Biesmeijer K, Smagghe G. 2015. Microsatellite analysis
581 of museum specimens reveals historical differences in genetic diversity between declining and more
582 stable *Bombus* species. *Plos One* 10: e0127870.

583 Martins IS, Proença V, Pereira HM. 2014. The unusual suspect: Land use is a key predictor of biodiversity
584 patterns in the Iberian Peninsula. *Acta Oecologica* 61: 41–50.

585 Miraldo A, Hewitt GM, Paulo OS, Emerson BC. 2011. Phylogeography and demographic history of *Lacerta*
586 *lepidota* in the Iberian Peninsula: multiple refugia, range expansions and secondary contact zones. *BMC*
587 *Evolutionary Biology* 11: 170.

588 Moreira AS, Horgan FG, Murray TE, Kakouli-Duarte T. 2015. Population genetic structure of *Bombus*
589 *terrestris* in Europe: Isolation and genetic differentiation of Irish and British populations. *Molecular*
590 *Ecology* 24: 3257–3268.

591 Murray TE, Coffey MF, Kehoe E, Horgan FG. 2013. Pathogen prevalence in commercially reared bumble
592 bees and evidence of spillover in conspecific populations. *Biological Conservation* 159: 269–276.

593 Newbold T, Hudson LN, Hill SLL, Contu S, Lysenko I, Senior RA, Börger L, Bennett DJ, Choimes A,
594 Collen B, Day J, De Palma A, Díaz S, Echeverria-Londoño S, Edgar MJ, Feldman A, Garon M,
595 Harrison MLK, Alhusseini T, Ingram DJ, Itescu Y, Kattge J, Kemp V, Kirkpatrick L, Kleyer M, Correia
596 DLP, Martin CD, Meiri S, Novosolov M, Pan Y, Phillips HRP, Purves DW, Robinson A, Simpson J,
597 Tuck SL, Weiher E, White HJ, Ewers RM, Mace GM, Scharlemann JPW, Purvis A. 2015. Global
598 effects of land use on local terrestrial biodiversity. *Nature* 520: 45–50.

599 Nunes VL, Mendes R, Marabuto E, Novais BM, Hertach T, Quartau JA, Seabra SG, Paulo OS, Simões PC.
600 2014. Conflicting patterns of DNA barcoding and taxonomy in the cicada genus *Tettigettalna* from
601 southern Europe (Hemiptera: Cicadidae). *Molecular Ecology Resources* 14: 27–38.

602 Penado A, Rebelo H, Goulson D. 2016. Spatial distribution modelling reveals climatically suitable areas for
603 bumblebees in undersampled parts of the Iberian Peninsula. *Insect Conservation and Diversity* 9: 391–
604 401.

605 Pina-Martins F, Silva DN, Fino J, Paulo OS. 2017. Structure_threader: An improved method for automation
606 and parallelization of programs Structure, fastStructure and MaverickK on multicore CPU systems.
607 *Molecular Ecology Resources* 17: e268–e274.

608 Pinto MA, Henriques D, Neto M, Guedes H, Muñoz I, Azevedo JC, De La Rúa P. 2013. Maternal diversity
609 patterns of Ibero-Atlantic populations reveal further complexity of Iberian honeybees. *Apidologie* 44:
610 430–439.

611 Plummer M, Best N, Cowles K, Vines K. 2006. CODA: Convergence diagnosis and output analysis for
612 MCMC. *R News* 6: 7–11.

613 Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010. Global pollinator declines:
614 Trends, impacts and drivers. *Trends in Ecology and Evolution* 25: 345–353.

615 Rasmont P, Coppée A, Michez D, De Meulemeester T. 2008. An overview of the *Bombus terrestris* (L. 1758)
616 subspecies (Hymenoptera: Apidae). *Annales de La Société Entomologique de France (N.S.)* 44: 243–
617 250.

618 Rasmont P, Franzén M, Lecocq T, Harpke A, Roberts SPM, Biesmeijer K, Cederberg B, Dvůrák L,
 619 Fitzpatrick Ú, Gonthier Y, Haubruge E, Mahé G, Manino A, Michez D, Neumayer J, Ødegaard F,
 620 Paukkunen J, Pawlikowski T, Potts SG, Reemer M, Settele J, Straka J, Schweiger O. 2015. Climatic
 621 risk and distribution atlas of European bumblebees. *BioRisk* 10: 1–236.

622 Rodrigues ASB, Silva SE, Marabuto E, Silva DN, Wilson MR, Thompson V, Yurtsever S, Halkka A, Borges
 623 PAV, Quartau JA, Paulo OS, Seabra SG. 2014. New mitochondrial and nuclear evidences support recent
 624 demographic expansion and an atypical phylogeographic pattern in the spittlebug *Philaenus spumarius*
 625 (Hemiptera, Aphrophoridae). *PLoS ONE* 9: e98375.

626 Seabra SG, Silva SE, Nunes VL, Sousa VC, Martins J, Marabuto E, Rodrigues ASB, Pina-Martins F,
 627 Laurentino TG, Rebelo MT, Figueiredo E, Paulo OS. 2019. Genomic signatures of introgression
 628 between commercial and native bumblebees, *Bombus terrestris*, in western Iberian Peninsula—
 629 Implications for conservation and trade regulation. *Evolutionary Applications* 12: 679–691.

630 Schmidt S, Schmid-Egger C, Morinière J, Haszprunar G, Hebert PD. 2015. DNA barcoding largely supports
 631 250 years of classical taxonomy: identifications for Central European bees (Hymenoptera, Apoidea
 632 *partim*). *Molecular Ecology Resources* 15: 985–1000.

633 Szulkin M, Gagnaire P-A, Bierne N, Charmantier A. 2016. Population genomic footprints of fine-scale
 634 differentiation between habitats in Mediterranean blue tits. *Molecular Ecology* 25: 542–558.

635 Trillo A, Montero-Castaño A, González-Varo JP, González-Moreno P, Ortiz-Sánchez FJ, Vilà M. 2019.
 636 Contrasting occurrence patterns of managed and native bumblebees in natural habitats across a
 637 greenhouse landscape gradient. *Agriculture, Ecosystems and Environment* 272: 230–236.

638 Verity R, Nichols RA. 2016. Estimating the number of subpopulations (K) in structured populations.
 639 *Genetics* 203: 1827–1835.

640 Velthuis HHW, van Doorn A. 2006. A century of advances in bumblebee domestication and the economic
 641 and environmental aspects of its commercialization for pollination. *Apidologie* 37: 421–451.

642 Vitalis R, Gautier M, Dawson KJ, Beaumont MA. 2014. Detecting and measuring selection from gene
 643 frequency data. *Genetics* 196: 799–817.

644 Walther-Hellwig K, Frankl R. 2000. Foraging habitats and foraging distances of bumblebees, *Bombus* spp.
645 (Hym., Apidae), in an agricultural landscape. *Journal of Applied Entomology* 124: 299–306.

646 Weingartner E, Wahlberg N, Nylin S. 2006. Speciation in *Pararge* (Satyrinae: Nymphalidae)
647 butterflies – North Africa is the source of ancestral populations of all *Pararge* species.
648 *Systematic Entomology* 31: 621-632.

649 Widmer A, Schmid-Hempel P, Estoup A, Scholls A. 1998. Population genetic structure and colonization
650 history of *Bombus terrestris* s.l. (Hymenoptera: Apidae) from the Canary Islands and Madeira. *Heredity*
651 81: 563–572.

652 Williams PH, An J, Brown MJF, Carolan JC, Goulson D, Huang J, Ito M. 2012a. Cryptic bumblebee
653 species: Consequences for conservation and the trade in greenhouse pollinators. *PLoS ONE* 7:
654 e32992.

655 Williams PH, Brown MJF, Carolan JC, An J, Goulson D, Aytekin AM, Best LR, Byvaltsev AM, Cederberg
656 B, Dawson R, Huang J, Ito M, Monfared A, Raina RH, Schmid-Hempel P, Sheffield CS, Šima P, Xie Z.
657 2012b. Unveiling cryptic species of the bumblebee subgenus *Bombus* s. str. worldwide with COI
658 barcodes (Hymenoptera: Apidae), *Systematics and Biodiversity* 10: 21-56.

659 Woodard SH, Lozier JD, Goulson D, Williams PH, Strange JP, Jha S. 2015. Molecular tools and bumble
660 bees: revealing hidden details of ecology and evolution in a model system. *Molecular Ecology* 24:
661 2916–2936.

662 **Figures and Tables**

664 **Figure 1.** Sampling locations of samples used in this study. A) Geographic distribution of each *B. terrestris*'
665 subspecies, according to Lecocq *et al.* (2016), here represented by different colours; geographic location of
666 collected samples of *B. t. terrestris*, *B. t. dalmatinus*, *B. t. audax* and *B. t. africanus*, as well as of COI
667 sequences (from previous studies of Coppée (2010), Williams *et al.* (2012a, b), Moreira *et al.* (2015) and
668 Schmidt *et al.* (2015)) downloaded from GenBank. B) Sampling locations of collected samples of *B. t.*
669 *lusitanicus* across the Iberian Peninsula (IP-NW: Iberian Peninsula, Northwest; IP-WE: Iberian Peninsula,

670 West; IP-SW: Iberian Peninsula, Southwest; IP-NO: Iberian Peninsula, North; IP-CE: Iberian Peninsula,
671 Centre; IP-SE: Iberian Peninsula, Southeast). B is a zoom-in of the black square in A. Sampling locations are
672 represented as follows: orange dots – samples used for COI analyses; yellow stars – samples used for both
673 COI and RAD analyses; black dots – COI sequences downloaded from GenBank.

674

675 **Figure 2.** Geographic distribution and frequency of mtDNA COI haplotypes. A) Map of haplotype
676 frequencies for *B. terrestris* across Europe and North Africa. B) Map of haplotype frequencies for each
677 Iberian geographic region. B is a zoom-in of the black square in A. C) Median-joining network representing
678 relationship among haplotypes, coloured by haplotype. Sequences from Coppée (2010), Williams *et al.*
679 (2012a, b), Moreira *et al.* (2015) and Schmidt *et al.* (2015) were also included. The size of the pie charts in
680 A, B and of the circles in C is in proportion to the haplotype frequencies. Each haplotype is represented by
681 the respective colour and designation (H1-H9 and H11-H17).

682

683 **Figure 3.** Principal components (PCA) and *Maverick* analyses of RADseq data. A) PCA of dataset
684 BT_OUT, comparing *B. terrestris* with other species; B) PCA of dataset BT_SSP comparing *B. t. lusitanicus*
685 samples with samples of other *B. terrestris*' subspecies; C) *Maverick* clustering plot of dataset BT_SSP, for
686 K=2; D) PCA of dataset BT_BTL comparing *B. t. lusitanicus* among Iberian geographical regions. In
687 *Maverick* results, estimated membership of each individual to each cluster are shown by vertical bars with
688 the clusters represented by different shades of grey.

689

690 **Table 1.** Results of the analyses of molecular variance (AMOVA) considering mitochondrial DNA COI data
691 and RADseq data.

692

693 **Table 2.** The sample size and diversity indices across *B. t. lusitanicus* samples from the Iberian Peninsula
694 considering mitochondrial DNA COI data and RADseq data, along with data for other populations of *B.*
695 *terrestris* or for other species (from Lozier, 2014 and Jackson *et al.*, 2018). Note that for RADseq markers, it
696 was also indicated the restriction enzyme used and the number of SNPs obtained.

697

698 **Supporting Information**

699

700 **Table S1.** List of samples used in this study: **Table S1a:** List of collected samples; **Table S1b:** List of
701 samples with sequences downloaded from GenBank.

702

703 **Table S2.** The sample size and diversity indices across *B. t. lusitanicus* samples by Iberian region
704 considering mitochondrial DNA COI data and RADseq dataset BT_BTL.

705

706 **Appendix 1.** Preliminary tests to identify optimal *Stacks* parameters: **Figure S1:** Results of parameter tests
707 for the *Stacks* module *pstacks*; **Figure S2:** results of parameter tests for the *Stacks* module *ctsacks* and
708 *sstacks*; **Figure S3:** results of parameter tests for the *Stacks* module *populations*.

709

710 **Appendix 2.** List of command line commands used for RADseq dataset analyses.

Figure 1




Figure 2




Figure 3




Table 1

DNA marker	Source of variation	Sum of squares	Variation components	Percentage of variation
COI	Within geographical regions	19.62	0.06	100.03
	Among geographical regions	0.27	-0.00002	-0.03
RAD seq	Within geographical regions	79454.05	1532.05	97.81
	Among geographical regions	9121.49	34.29	2.19

Table 2

Species	Geographic region	Reference	Sample size	DNA marker	h (n° of haplotypes)	π	π_{SNP}	H_o	H_e
<i>B. t. lusitanicus</i>	Iberian Peninsula	this study	208	COI	0.08470 (7 hap.)	0.00018	-	-	-
<i>Bombus terrestris</i>	Europe (continental)	this study	340	COI	0.09740 (10 hap.)	0.00034	-	-	-
<i>Bombus terrestris</i>	Europe (continental + islands) & north Africa	this study	441	COI	0.39900 (16 hap.)	0.00209	-	-	-
<i>B. t. lusitanicus</i>	Iberian Peninsula	this study	31 ♀	RAD seq, PstI (10,938 SNPs)	-	-	0.278	0.233	0.277
<i>Bombus impatiens</i>	eastern U.S.A.	Lozier <i>et al.</i> (2014)	24 ♀	RAD seq, SgrAI (2,387 – 9,148 SNPs)	-	-	0.136 – 0.289	n.d.	n.d.
<i>Bombus pensylvanicus</i>	eastern U.S.A.	Lozier <i>et al.</i> (2014)	24 ♀	RAD seq, SgrAI (3,240 – 9,376 SNPs)	-	-	0.135 – 0.276	n.d.	n.d.
<i>Bombus bifarius</i>	mountain regions of western U.S.A.	Jackson <i>et al.</i> (2018)	383 ♀	RAD seq, PstI (598 – 37,474 SNPs)	-	-	0.122 – 0.140	n.d.	n.d.
<i>Bombus vosnesenskii</i>	mountain regions of western U.S.A.	Jackson <i>et al.</i> (2018)	587 ♀	RAD seq, PstI (356 – 18,700 SNPs)	-	-	0.105 – 0.116	n.d.	n.d.

h , haplotype diversity; π , nucleotide diversity; π_{SNP} , per-SNP nucleotide diversity; H_o , mean per-individual observed heterozygosity; H_e , mean per-individual expected heterozygosity. “-” indicates “non-applicable”; “n.d.” indicates “no data”.

Table S1a. List of collected samples with information concerning: sampling (taxa, sex, country, number of samples, geographic groups, origin, date of collection and respective collectors); outputs of mitochondrial COI analyses (haplotypes: mapped reads, coverage, datasets at which belongs and missing data).

B. t. lusitanicus

Sample Code	Taxa	Sex	n	Country / Island	Geographic region	Sampling Location	Latitude	Longitude	Date of collection	Collector(s)	mtDNA analysis	COI haplotype	GenBank accession no.
BOM_GEH_002	<i>B. t. lusitanicus</i>	F	1	Portugal	IP-NW Iberian Peninsula, northwest (n=24)	Montalegre	41.72247	-7.68883	03/07/2017	S.E. Silva & R. Mendes	x	H1	MN652675
BOM_GEH_003	<i>B. t. lusitanicus</i>	F	1	Portugal		Montalegre	41.72247	-7.68883	03/07/2017	S.E. Silva & R. Mendes	x	H1	MN652676
BOM_GEH_006	<i>B. t. lusitanicus</i>	F	1	Portugal		Montalegre, Parafita	41.75589	-7.84843	03/07/2017	S.E. Silva & R. Mendes	x	H1	MN652677
BOM_GEH_009	<i>B. t. lusitanicus</i>	F	1	Portugal		Montalegre, Parafita	41.75589	-7.84843	03/07/2017	S.E. Silva & R. Mendes	x	H1	MN652678
BOM_GEH_028	<i>B. t. lusitanicus</i>	F	1	Portugal		P.N. Peneda-Gerês, Vilar da Veiga	41.73505	-8.159281	15/05/2014	T.G. Laurentino & O.S. Paulo	x	H1	MN652679
BTL_064	<i>B. t. lusitanicus</i>	F	1	Portugal		P.N. Peneda-Gerês, Vilar da Veiga	41.73505	-8.159281	15/05/2014	T.G. Laurentino & O.S. Paulo	x	H1	MN652680
BTL_065	<i>B. t. lusitanicus</i>	F	1	Portugal		P.N. Peneda-Gerês, Vilar da Veiga	41.73505	-8.159281	15/05/2014	T.G. Laurentino & O.S. Paulo	x	H1	MN652681
BTL_060	<i>B. t. lusitanicus</i>	F	1	Portugal		P.N. Montesinho, Pinheiro Novo	41.935667	-7.114667	13/05/2014	T.G. Laurentino & O.S. Paulo	x	H1	MN652682
BTL_061	<i>B. t. lusitanicus</i>	F	1	Portugal		P.N. Montesinho, Pinheiro Novo	41.935667	-7.114667	13/05/2014	T.G. Laurentino & O.S. Paulo	x	H1	MN652683
BTL_062	<i>B. t. lusitanicus</i>	F	1	Portugal		P.N. Montesinho, Pinheiro Novo	41.937944	-7.108444	13/05/2014	T.G. Laurentino & O.S. Paulo	x	H1	MN652684
BOM_GEL_002	<i>B. t. lusitanicus</i>	F	1	Spain		Pontevedra, Pazos de Borbén	42.277028	-8.525778	02/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652685
BOM_GEL_010	<i>B. t. lusitanicus</i>	F	1	Spain		Pontevedra, Pazos de Borbén	42.302028	-8.546667	02/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652686
BOM_GEL_016	<i>B. t. lusitanicus</i>	F	1	Spain		Pontevedra, Soutomaior	42.331472	-8.563556	02/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652687
BOM_GEL_021	<i>B. t. lusitanicus</i>	F	1	Spain		Pontevedra, Soutomaior	42.308389	-8.5495	02/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652688
BOM_GEL_026	<i>B. t. lusitanicus</i>	F	1	Spain		Pontevedra, Soutomaior	42.339139	-8.471333	02/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652689
BOM_MLL_001	<i>B. t. lusitanicus</i>	F	1	Spain		Monforte de Lemos, Canabal	42.48316	-7.59038	04/07/2017	S.E. Silva & R. Mendes	x	H1	MN652690
BOM_MLL_003	<i>B. t. lusitanicus</i>	F	1	Spain		Monforte de Lemos, Pantón	42.48619	-7.59861	04/07/2017	S.E. Silva & R. Mendes	x	H1	MN652691
BOM_MLL_010	<i>B. t. lusitanicus</i>	F	1	Spain		Monforte de Lemos, Pantón	42.51736	-7.67458	04/07/2017	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652692
BOM_MLL_011	<i>B. t. lusitanicus</i>	F	1	Spain		Lugo, Monforte de Lemos	42.54425	-7.603667	03/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652693
BOM_MLL_026	<i>B. t. lusitanicus</i>	F	1	Spain		Lugo, Monforte de Lemos	42.549833	-7.612667	03/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652694
BTL_075	<i>B. t. lusitanicus</i>	F	1	Spain		Ourense, Serra da Enciña da Lastra	42.486833	-6.860333	12/05/2014	E. Marabuto	x	H1	MN652695
BTL_076	<i>B. t. lusitanicus</i>	F	1	Spain		Ourense, Serra da Enciña da Lastra	42.486833	-6.860333	12/05/2014	E. Marabuto	x	H1	MN652696
BOM_MLH_010	<i>B. t. lusitanicus</i>	F	1	Spain		Ourense, A Veiga	42.232389	-6.941028	03/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652697
BOM_MLH_016	<i>B. t. lusitanicus</i>	F	1	Spain		Ourense, Serra da Enciña da Lastra	42.486833	-6.860333	12/05/2014	E. Marabuto	x	H4	MN652698
BOM_SMH_003	<i>B. t. lusitanicus</i>	F	1	Portugal	IP-WE Iberian Peninsula, west (n=56)	Serra de São Mamede	39.310361	-7.380139	23/04/2017	S.E. Silva	x	H1	MN652699
BOM_SMH_004	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra de São Mamede	39.310361	-7.380139	23/04/2017	S.E. Silva	x	H1	MN652700
BOM_SMH_005	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra de São Mamede	39.308694	-7.393694	23/04/2017	S.E. Silva	x	H1	MN652701
BOM_SMH_006	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra de São Mamede	39.308694	-7.393694	23/04/2017	S.E. Silva	x	H1	MN652702
BOM_SMH_013	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra de São Mamede	39.312944	-7.359917	19/06/2016	S.E. Silva	x	H1	MN652703
BOM_SML_001	<i>B. t. lusitanicus</i>	F	1	Portugal		Portalegre, Gafete	39.412861	-7.651167	23/04/2017	S.E. Silva	x	H1	MN652704
BTL_306	<i>B. t. lusitanicus</i>	F	1	Portugal		Abrantes, Mouriscas	39.508733	-8.103575	08/04/2012	E. Figueiredo	x	H1	MN652705
BTL_048	<i>B. t. lusitanicus</i>	F	1	Portugal		Coimbra, Larçã	40.327167	-8.409417	06/05/2014	E. Marabuto	x	H1	MN652706
BTL_050	<i>B. t. lusitanicus</i>	F	1	Portugal		Coimbra, Rabaçal	40.031833	-8.435833	05/05/2014	E. Marabuto	x	H1	MN652707
BOM_CML_001	<i>B. t. lusitanicus</i>	F	1	Portugal		Mealhada, Pampilhosa do Botão	40.322861	-8.441917	21/05/2017	S.E. Silva	x	H1	MN652708
BOM_CML_002	<i>B. t. lusitanicus</i>	F	1	Portugal		Mealhada, Pampilhosa do Botão	40.331306	-8.423778	21/05/2017	S.E. Silva	x	H1	MN652709
BOM_CML_003	<i>B. t. lusitanicus</i>	F	1	Portugal		Mealhada, Pampilhosa do Botão	40.331306	-8.423778	21/05/2017	S.E. Silva	x	H1	MN652710
BOM_CML_006	<i>B. t. lusitanicus</i>	F	1	Portugal		Mealhada, Pampilhosa do Botão	40.328694	-8.40275	21/05/2017	S.E. Silva	x	H1	MN652711
BOM_CML_007	<i>B. t. lusitanicus</i>	F	1	Portugal		Mealhada, Pampilhosa do Botão	40.328694	-8.40275	21/05/2017	S.E. Silva	x	H1	MN652712
BOM_CMH_006	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra do Caramulo	40.573694	-8.157056	21/05/2017	S.E. Silva	x	H1	MN652713
BOM_CMH_009	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra do Caramulo	40.554528	-8.195139	21/05/2017	S.E. Silva	x	H1	MN652714
BOM_CMH_011	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra do Caramulo	40.554528	-8.195139	21/05/2017	S.E. Silva	x	H1	MN652715
BOM_CMH_012	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra do Caramulo	40.548806	-8.202083	21/05/2017	S.E. Silva	x	H1	MN652716
BOM_CMH_015	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra do Caramulo	40.554167	-8.198556	01/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652717
BTL_316	<i>B. t. lusitanicus</i>	F	1	Portugal		Oliveira de Frades, Reigoso	40.67725	-8.274169	16/06/2013	E. Figueiredo	x	H1	MN652718
BOM_SEL_001	<i>B. t. lusitanicus</i>	F	1	Portugal		Penalva do Castelo	40.644139	-7.713361	04/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652719
BOM_SEL_002	<i>B. t. lusitanicus</i>	F	1	Portugal		Penalva do Castelo	40.644139	-7.713361	04/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H2	MN652720
BOM_SEL_008	<i>B. t. lusitanicus</i>	F	1	Portugal		Penalva do Castelo	40.663583	-7.683472	04/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652721
BOM_SEL_013	<i>B. t. lusitanicus</i>	F	1	Portugal		Penalva do Castelo	40.671056	-7.641194	04/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652722
BOM_SEL_026	<i>B. t. lusitanicus</i>	F	1	Portugal		Penalva do Castelo	40.665444	-7.657806	04/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652723
BTL_203	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra da Estrela	40.431731	-7.51575	11/06/2014	T.G. Laurentino	x	H1	MN652724
BTL_213	<i>B. t. lusitanicus</i>	F	1	Portugal		Serra da Estrela	40.365167	-7.642167	11/06/2014	E. Marabuto	x	H1	MN652725

BTL_217	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_220	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SEH_002	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SEH_007	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SEH_010	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SEH_027	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SEH_029	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_001	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_297	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_021	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_037	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_059	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_324	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_278	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_281	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_282	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_283	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_288	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAL_002	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAL_003	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAL_004	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAL_005	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAL_018	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_201	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAH_006	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAH_008	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAH_009	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAH_013	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_SAH_014	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_MOH_001	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_MOH_002	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_MOH_008	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_200	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_356	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_357	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_358	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_346	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_347	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_349	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_MOL_001	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_MOL_005	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_MOL_007	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_CAH_002	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_CAH_004	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_CAH_006	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_CAL_001	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_CAL_003	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_CAL_009	<i>B. t. lusitanicus</i>	F	1	Portugal
BTL_235	<i>B. t. lusitanicus</i>	F	1	Portugal
BOM_PIL_001	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_PIL_008	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_PIL_009	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_PIL_011	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_PIL_012	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_PIH_021	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_NAV_002	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_NAV_004	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_077	<i>B. t. lusitanicus</i>	?	1	Spain
BTL_078	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_092	<i>B. t. lusitanicus</i>	F	1	Spain

IP-SW

Iberian Peninsula, southwest

(n=20)

IP-NO

Iberian Peninsula, north

(n=28)

Serra da Estrela	40.410833	-7.671833	11/06/2014	E. Marabuto	x	H1	MN652726
Serra da Estrela	40.365167	-7.642167	11/06/2014	E. Marabuto	x	H1	MN652727
Serra da Estrela	40.408528	-7.663583	04/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652728
Serra da Estrela	40.406306	-7.566139	04/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652729
Serra da Estrela	40.398917	-7.552556	04/07/2016	S.E. Silva, R. Mendes & M.J. Does	x	H1	MN652730
Serra da Estrela	40.410833	-7.671833	11/06/2014	E. Marabuto	x	H1	MN652731
Serra da Estrela	40.365167	-7.642167	11/06/2014	E. Marabuto	x	H1	MN652732
Oeiras, Lage	38.709267	-9.311717	10/04/2014	E. Marabuto	x	H1	MHO18622
Oeiras	38.690789	-9.313989	25/07/2014	V.L. Nunes	x	H1	MHO18657
Cascais	38.743	-9.4307	27/04/2014	E. Marabuto	x	H1	MHO18623
Lisboa, near the airport	38.768117	-9.148017	01/05/2014	V.L. Nunes	x	H1	MHO18626
Lisboa, near the airport	38.768117	-9.148017	13/05/2014	V.L. Nunes	x	H1	MHO18627
Lisboa, Tapada da Ajuda	38.707639	-9.182306	13/04/2014	E. Figueiredo	x	H1	MHO18660
Bombarral, Portela	39.244861	-9.134778	17/07/2014	O.S. Paulo, S.E. Silva & A.S. Rodrigues	x	H1	MHO18652
Cadaval	39.240472	-9.099444	17/07/2014	O.S. Paulo, S.E. Silva & A.S. Rodrigues	x	H1	MHO18653
Cadaval, Vilar	39.188861	-9.112944	17/07/2014	O.S. Paulo, S.E. Silva & A.S. Rodrigues	x	H1	MHO18654
Alenquer, Vila Verde dos Francos	39.153389	-9.112583	17/07/2014	O.S. Paulo, S.E. Silva & A.S. Rodrigues	x	H1	MHO18655
Alenquer, Vila Verde dos Francos	39.153389	-9.112583	17/07/2014	O.S. Paulo, S.E. Silva & A.S. Rodrigues	x	H1	MHO18656
Alenquer, Abrigada	39.136833	-9.019722	16/04/2017	S.E. Silva	x	H1	MN652733
Alenquer, Abrigada	39.136833	-9.019722	16/04/2017	S.E. Silva	x	H1	MN652734
Alenquer, Abrigada	39.143333	-9.038556	16/04/2017	S.E. Silva	x	H1	MN652735
Alenquer, Abrigada	39.143333	-9.038556	16/04/2017	S.E. Silva	x	H1	MN652736
Alenquer, Ota	39.10375	-9.0115	26/06/2016	S.E. Silva	x	H1	MN652737
Foz do Arelho	39.429139	-9.223472	11/06/2014	B. Costa	x	H1	MHO18630
Serra de Aire e Candeeiros	39.457528	-8.900222	26/06/2016	S.E. Silva	x	H2	MN652738
Serra de Aire e Candeeiros	39.471139	-8.904333	26/06/2016	S.E. Silva	x	H1	MN652739
Serra de Aire e Candeeiros	39.471139	-8.904333	26/06/2016	S.E. Silva	x	H1	MN652740
Serra de Aire e Candeeiros	39.478861	-8.888917	26/06/2016	S.E. Silva	x	H1	MN652741
Serra de Aire e Candeeiros	39.503083	-8.872111	26/06/2016	S.E. Silva	x	H1	MN652742
Serra de Monchique	37.321593	-8.595798	18/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652743
Serra de Monchique	37.321593	-8.595798	18/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652744
Serra de Monchique	37.317333	-8.583278	18/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652745
Silves, Armação de Pêra	37.102358	-8.368486	08/06/2014	V.L. Nunes	x	H1	MN652746
Sines, Porto Covo	37.854286	-8.793353	16/06/2014	E. Figueiredo	x	H1	MHO18670
Sines, Porto Covo	37.854286	-8.793353	16/06/2014	E. Figueiredo	x	H3	MHO18671
Sines, Porto Covo	37.854286	-8.793353	16/06/2014	E. Figueiredo	x	H1	MHO18672
Odemira, Zambujeira do Mar	37.534159	-8.785925	16/06/2014	E. Figueiredo	x	H1	MN652747
Odemira, Zambujeira do Mar	37.534159	-8.785925	16/06/2014	E. Figueiredo	x	H1	MN652748
Odemira, Zambujeira do Mar	37.534159	-8.785925	16/06/2014	E. Figueiredo	x	H1	MN652749
Odemira, Luzianes	37.5911472	-8.488128	18/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652750
Odemira, Azinhaga do Calvário	37.514778	-8.476028	18/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652751
Odemira, Santa Clara a Velha	37.513111	-8.471389	18/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652752
Serra do Caldeirão, Alcaria do Cume	37.204217	-7.687183	22/05/2016	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652753
Serra do Caldeirão, Alcaria do Cume	37.219033	-7.705183	22/05/2016	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652754
Serra do Caldeirão, Alcaria do Cume	37.244667	-7.742817	22/05/2016	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652755
Mértola, Corte de Gafo de Cima	37.7194141	-7.7067368	19/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652756
Almodôvar	37.5124	-8.057394	19/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652757
Almodôvar	37.5127963	-8.0692471	19/03/2017	S.E. Silva, A.S.B. Rodrigues & O.S. Paulo	x	H1	MN652758
Beja, Beringel	38.055417	-7.9973	08/07/2014	R. Mendes	x	H1	MN652759
Navarra, Caparrosa	42.300444	-1.650056	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652760
Navarra, Mélida	42.360944	-1.546083	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652761
Navarra, Mélida	42.360944	-1.546083	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652762
Navarra, Carcastillo	42.371722	-1.457694	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652763
Navarra, Carcastillo	42.371722	-1.457694	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652764
Navarra, Abaurregaina	42.902528	-1.187833	12/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652765
Navarra, Pamplona	42.797	-1.627778	29/06/2014	E. Figueiredo	x	H1	MN652766
Navarra, Pamplona	42.797	-1.627778	29/06/2014	E. Figueiredo	x	H1	MN652767
Cantabria, Vega de Liébana	43.120833	-4.626	14/05/2014	E. Marabuto	x	H1	MN652768
Cantabria, Vega de Liébana	43.120833	-4.626	14/05/2014	E. Marabuto	x	H1	MN652769
Cantabria, Vega de Liébana	43.086333	-4.7245	14/05/2014	E. Marabuto	x	H1	MN652770

BTL_094	<i>B. t. lusitanicus</i>	F	1	Spain	IP-CE Iberian Peninsula, centre (n=32)	Cantabria, Vega de Liébana	43.086333	-4.7245	14/05/2014	E. Marabuto	x	H1	MN652771
BTL_099	<i>B. t. lusitanicus</i>	F	1	Spain		Cantabria, Vega de Liébana	43.086333	-4.7245	14/05/2014	E. Marabuto	x	H1	MN652772
BOM_CTL_002	<i>B. t. lusitanicus</i>	F	1	Spain		Navarra, Mendaza	42.684667	-2.270778	12/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652773
BOM_CTL_003	<i>B. t. lusitanicus</i>	F	1	Spain		Navarra, Mendaza	42.684667	-2.270778	12/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652774
BOM_CTL_010	<i>B. t. lusitanicus</i>	F	1	Spain		Álava, Bernedo	42.627278	-2.504111	12/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652775
BOM_CTH_002	<i>B. t. lusitanicus</i>	F	1	Spain		Burgos, Merindad de Cuesta	42.858028	-3.4275	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652776
BOM_CTH_003	<i>B. t. lusitanicus</i>	F	1	Spain		Burgos, Merindad de Cuesta	42.858028	-3.4275	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652777
BOM_PEH_001	<i>B. t. lusitanicus</i>	F	1	Spain		Cantabria, Allende	43.21725	-4.593722	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652778
BOM_PEH_002	<i>B. t. lusitanicus</i>	F	1	Spain		Cantabria, Allende	43.21725	-4.593722	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652779
BOM_PEH_003	<i>B. t. lusitanicus</i>	F	1	Spain		Cantabria, Allende	43.21725	-4.593722	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652780
BOM_PEH_010	<i>B. t. lusitanicus</i>	F	1	Spain		Léon, Boca de Huérgano	42.984167	-4.907944	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652781
BOM_PEH_013	<i>B. t. lusitanicus</i>	F	1	Spain		Léon, Boca de Huérgano	42.984167	-4.907944	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652782
BOM_PEL_002	<i>B. t. lusitanicus</i>	F	1	Spain		Léon, Cistierna	42.748556	-5.140139	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652783
BOM_PEL_006	<i>B. t. lusitanicus</i>	F	1	Spain		Léon, Cubillas de Rueda	42.7285	-5.1455	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652784
BOM_PEL_007	<i>B. t. lusitanicus</i>	F	1	Spain		Léon, Cubillas de Rueda	42.7285	-5.1455	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H2	MN652785
BOM_PEL_010	<i>B. t. lusitanicus</i>	F	1	Spain		Léon, Cubillas de Rueda	42.676	-5.169167	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652786
BOM_PEL_011	<i>B. t. lusitanicus</i>	F	1	Spain		Léon, Cubillas de Rueda	42.676	-5.169167	13/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652787
BOM_GDH_001	<i>B. t. lusitanicus</i>	F	1	Spain		Ávila, Sierra de Gredos, Cuevas de Valle	40.309167	-5.016778	10/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652788
BOM_TOL_001	<i>B. t. lusitanicus</i>	F	1	Spain		Ciudad Real, Santa Quiteria	39.257099	-4.3604809	03/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652789
BOM_TOL_002	<i>B. t. lusitanicus</i>	F	1	Spain		Ciudad Real, Santa Quiteria	39.257099	-4.3604809	03/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652790
BOM_TOH_010	<i>B. t. lusitanicus</i>	F	1	Spain		Toledo, Montes de Toledo, Hontanar	39.586946	-4.506851	03/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652791
BOM_TOH_011	<i>B. t. lusitanicus</i>	F	1	Spain		Toledo, Montes de Toledo, Hontanar	39.586946	-4.506851	03/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652792
BOM_TOH_015	<i>B. t. lusitanicus</i>	F	1	Spain		Toledo, Montes de Toledo, Hontanar	39.5865000001	-4.507	15/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652793
BOM_TOH_025	<i>B. t. lusitanicus</i>	F	1	Spain		Toledo, Montes de Toledo, Hontanar	39.5752237423	-4.5410353884	15/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652794
BOM_TOH_028	<i>B. t. lusitanicus</i>	F	1	Spain		Toledo, Montes de Toledo, Valdeazores	39.4500492018	-4.6827687569	15/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652795
BOM_GUH_003	<i>B. t. lusitanicus</i>	F	1	Spain		Cáceres, Sierra de Guadalupe, Guadalupe	39.4615341	-5.3295201	04/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652796
BOM_GUH_004	<i>B. t. lusitanicus</i>	F	1	Spain		Cáceres, Sierra de Guadalupe, Cañamero	39.4725	-5.362778	04/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652797
BOM_GUH_006	<i>B. t. lusitanicus</i>	F	1	Spain		Cáceres, Sierra de Guadalupe, Cañamero	39.476167	-5.36725	04/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652798
BOM_GUH_016	<i>B. t. lusitanicus</i>	F	1	Spain		Cáceres, Sierra de Guadalupe, Alía	39.500917	-5.345	04/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652799
BOM_GUH_017	<i>B. t. lusitanicus</i>	F	1	Spain		Cáceres, Sierra de Guadalupe, Alía	39.500917	-5.345	04/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652800
BTL_150	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Loeches	40.371944	-3.379861	26/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652801
BTL_144	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Rascafría	40.878583	-3.847806	26/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652802
BTL_145	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Rascafría	40.878583	-3.847806	26/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652803
BTL_148	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Rascafría	40.878583	-3.847806	26/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652804
BTL_301	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Real Sitio de San Ildefonso	40.900481	-4.009089	13/08/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652805
BOM_GRH_002	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Real Sitio de San Ildefonso	40.861	-4.027333	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652806
BOM_GRH_004	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Real Sitio de San Ildefonso	40.861	-4.027333	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652807
BOM_GRH_008	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Rascafría	40.872	-3.885	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652808
BOM_GRH_009	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Rascafría	40.872	-3.885	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652809
BOM_GRH_023	<i>B. t. lusitanicus</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Rascafría	40.878583	-3.847806	26/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652810
BOM_GRL_002	<i>B. t. lusitanicus</i>	F	1	Spain		Segovia, Cerezo de Arriba	41.245	-3.548722	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652811
BOM_GRL_003	<i>B. t. lusitanicus</i>	F	1	Spain		Segovia, Ribota	41.358972	-3.44675	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652812
BOM_GRL_004	<i>B. t. lusitanicus</i>	F	1	Spain		Segovia, Ribota	41.358972	-3.44675	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652813
BOM_GRL_009	<i>B. t. lusitanicus</i>	F	1	Spain		Soria, Valdeavillo	41.649472	-2.891194	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652814
BOM_GRL_010	<i>B. t. lusitanicus</i>	F	1	Spain		Soria, Villaciervos	41.752611	-2.651472	11/06/2017	S.E. Silva, R. Mendes & A.S.B. Rodrigues	x	H1	MN652815
BTL_086	<i>B. t. lusitanicus</i>	F	1	Spain	IP-SE Iberian Peninsula, southeast (n=38)	Valladolid, Castronuevo de Esgueva	41.686667	-4.597	16/05/2014	E. Marabuto	x	H1	MN652816
BTL_089	<i>B. t. lusitanicus</i>	F	1	Spain		Valladolid, Castronuevo de Esgueva	41.686667	-4.597	16/05/2014	E. Marabuto	x	H1	MN652817
BTL_090	<i>B. t. lusitanicus</i>	F	1	Spain		Valladolid, Castronuevo de Esgueva	41.686667	-4.597	16/05/2014	E. Marabuto	x	H1	MN652818
BTL_097	<i>B. t. lusitanicus</i>	F	1	Spain		Valladolid, Castronuevo de Esgueva	41.686667	-4.597	16/05/2014	E. Marabuto	x	H1	MN652819
BOM_ESH_003	<i>B. t. lusitanicus</i>	F	1	Spain		Murcia, Sierra Espuña	37.8627000005	-1.55323	12/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652820
BOM_ESH_010	<i>B. t. lusitanicus</i>	F	1	Spain		Murcia, Sierra Espuña	37.8650027781	-1.5713306989	12/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652821
BOM_ESH_013	<i>B. t. lusitanicus</i>	F	1	Spain		Murcia, Sierra Espuña	37.85508	-1.568043	11/07/2017	S.E. Silva & R. Mendes	x	H3	MN652822
BTL_115	<i>B. t. lusitanicus</i>	F	1	Spain		Granada, Puebla de Don Fadrique	37.953222	-2.408	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652823
BTL_118	<i>B. t. lusitanicus</i>	F	1	Spain		Granada, Puebla de Don Fadrique	38.042361	-2.472611	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652824
BTL_121	<i>B. t. lusitanicus</i>	F	1	Spain		Granada, Puebla de Don Fadrique	37.953222	-2.408	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652825
BTL_123	<i>B. t. lusitanicus</i>	F	1	Spain		Granada, Puebla de Don Fadrique	38.064417	-2.5215	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652826
BTL_124	<i>B. t. lusitanicus</i>	F	1	Spain		Granada, Puebla de Don Fadrique	38.042361	-2.472611	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652827
BTL_125	<i>B. t. lusitanicus</i>	F	1	Spain		Granada, Puebla de Don Fadrique	37.953222	-2.408	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H5	MN652828
BTL_109	<i>B. t. lusitanicus</i>	F	1	Spain		Almería, Sierra María-Los Vélez	37.694417	-2.174639	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652829
BTL_110	<i>B. t. lusitanicus</i>	F	1	Spain		Almería, Sierra María-Los Vélez	37.694417	-2.174639	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652830

BTL_112	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_136	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_138	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_141	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ALH_001	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ALH_003	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ALH_009	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ALH_016	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROL_001	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROL_002	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROL_008	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROL_009	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROH_001	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROH_002	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROH_003	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROH_004	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_ROH_014	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_185	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_191	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_197	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_168	<i>B. t. lusitanicus</i>	F	1	Spain
BTL_180	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_NEH_006	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_NEH_007	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_NEH_011	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_NEH_013	<i>B. t. lusitanicus</i>	F	1	Spain
BOM_NEH_038	<i>B. t. lusitanicus</i>	F	1	Spain

Almería, Sierra María-Los Vélez	37.694417	-2.174639	28/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652831
Alicante, Serra de Serrella-Barranc Fort	38.697333	-0.309361	27/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652832
Alicante, Serra de Serrella-Barranc Fort	38.697333	-0.309361	27/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652833
Alicante, Serra de Serrella-Barranc Fort	38.697333	-0.309361	27/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652834
Alicante, Ibi	38.6356000015	-0.5228	13/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H6	MN652835
Alicante, Serra de Serrella-Barranc Fort	38.7025000015	-0.3031000003	13/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652836
Alicante, Serra de Serrella-Barranc Fort	38.697333	-0.309361	27/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652837
Alicante, Serra de Serrella-Barranc Fort	38.697333	-0.309361	27/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652838
Sevilla, Montellano	36.993333	-5.569361	01/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652839
Sevilla, Montellano	36.993333	-5.569361	01/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652840
Sevilla, Montellano	36.984194	-5.545111	01/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652841
Sevilla, Montellano	36.984194	-5.545111	01/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652842
Málaga, Serranía de Ronda, Pujerra	36.612167	-5.140722	01/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652843
Málaga, Serranía de Ronda, Pujerra	36.612167	-5.140722	01/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H7	MN652844
Málaga, Serranía de Ronda, Pujerra	36.612778	-5.139194	01/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652845
Málaga, Serranía de Ronda, Pujerra	36.612778	-5.139194	01/04/2017	S.E. Silva, R. Mendes & O.S. Paulo	x	H1	MN652846
Málaga, El Burgo	36.8001999999	-4.9138	08/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652847
Granada, Sierra Nevada, El Molinillo	37.310833	-3.413889	29/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652848
Granada, Sierra Nevada, Alfacar	37.258889	-3.549167	29/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652849
Granada, Sierra Nevada, Alfacar	37.258889	-3.549167	29/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652850
Granada, Sierra Nevada, Monachil	37.139167	-3.467778	29/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652851
Granada, Sierra Nevada, Monachil	37.139167	-3.467778	29/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652852
Granada, Sierra Nevada, Alfacar	37.2592	-3.54825	09/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652853
Granada, Sierra Nevada, Alfacar	37.2592	-3.54825	09/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652854
Granada, Sierra Nevada, Alfacar	37.2660500741	-3.5265	09/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652855
Granada, Sierra Nevada, Monachil	37.1130500004	-3.4397	09/06/2016	S.E. Silva, R. Mendes & E. Marabuto	x	H1	MN652856
Granada, Sierra Nevada, Alfacar	37.258889	-3.549167	29/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H1	MN652857

Other Subspecies

Sample Code	Taxa	Sex	n	Country / Island	Geographic region	Sampling Location	Latitude	Longitude	Date of collection	Collector(s)	mtDNA analysis	COI haplotype	GenBank accession no.
BOM_SWI_003	<i>B. t. terrestris</i>	F	1	Switzerland	Switzerland	Basel	--	--	08/2016	T.G. Laurentino	x	H2	MN652858
BT_046	<i>B. t. terrestris</i>	F	1	Germany	Germany	Rothenburg ob der Tauber	49.374933	10.17695	08/05/2014	V.L. Nunes	x	H1	MN652859
BT_029	<i>B. t. terrestris</i>	M	1	France	France	Sainte-Anastasie-sur-Issole	43.3555	6.141	23/04/2014	E. Marabuto	x	H1	MN652860
BOM_TUR_001	<i>B. t. dalmatinus</i>	F	1	Turkey	Turkey	Tekirdag, Ganos Mountains	--	--	06/2016	S. Yurtsever	x	H1	MN652861
BTX_395	<i>B. t. audax</i>	F	1	Great Britain	England	P. Hill near Oxford	--	--	--	T.G. Laurentino	x	H8	MN652862
BTA_258	<i>B. t. africanus</i>	F	1	Morocco	Morocco (n=4)	Chefchaouen	35.182969	-5.222889	15/07/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H9	MN652863
BTA_261	<i>B. t. africanus</i>	F	1	Morocco		Chefchaouen	35.182969	-5.222889	15/07/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H9	MN652864
BTA_262	<i>B. t. africanus</i>	F	1	Morocco		Azrou	33.406333	-5.2035	16/07/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H9	MN652865
BTA_104	<i>B. t. africanus</i>	F	1	Morocco		Chefchaouen	35.104	-5.139417	18/07/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	H9	MN652866

Outgroup

Sample Code	Taxa	Sex	n	Country / Island	Geographic region	Sampling Location	Latitude	Longitude	Date of collection	Collector(s)	mtDNA analysis	COI haplotype	GenBank accession no.
BHO_067	<i>B. hortorum</i>	F	1	Portugal	Outgroup (n=13)	P.N. Peneda-Gerês, Vilar da Veiga	41.73505	-8.159281	14/05/2014	T.G. Laurentino & O.S. Paulo	x	-	MN652867
BLA_205	<i>B. lapidarius</i>	F	1	Portugal		Serra da Estrela	40.427972	-7.654022	10/06/2014	T.G. Laurentino	x	-	MN652868
BLA_212	<i>B. lapidarius</i>	F	1	Portugal		Serra da Estrela	40.462928	-7.513517	11/06/2014	T.G. Laurentino	x	-	MN652869
BLA_069	<i>B. lapidarius</i>	F	1	Spain		Cantabria, Vega de Liébana	43.086333	-4.7245	14/05/2014	E. Marabuto	x	-	MN652870
BTL_084	<i>B. lucorum</i>	F	1	Spain		Burgos, Barbadillo de Herreros	42.136667	-3.1935	15/05/2014	E. Marabuto	x	-	MHO18628
BTL_103	<i>B. lucorum</i>	F	1	Spain		Léon, Sierra de Ancares	42.871333	-6.452083	12/05/2014	E. Marabuto	x	-	MN652871
BTL_146	<i>B. lucorum</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Rascafría	40.878583	-3.847806	26/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	-	MHO18629
BPA_229	<i>B. pascuorum</i>	F	1	Portugal		Mafra, Sobral da Abelheira	--	--	28/06/2014	O.S. Paulo, S.E. Silva & A.S. Rodrigues	x	-	MN652872
BPA_068	<i>B. pascuorum</i>	F	1	Spain		Cantabria, Vega de Liébana	43.086333	-4.7245	14/05/2014	E. Marabuto	x	-	MN652873
BPA_133	<i>B. pascuorum</i>	F	1	Spain		Alicante, Serra de Serrella-Barranc Fort	38.690556	-0.287111	27/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	-	MN652874
BPA_152	<i>B. pascuorum</i>	F	1	Spain		Madrid, Sierra de Guadarrama, Miraflores de La Sierra	40.800278	-3.756667	26/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	-	MN652875

BHO_223	<i>B. ruderatus</i>	F	1	Portugal	Torres Vedras, Coutada	--	--	28/06/2014	O.S. Paulo, S.E. Silva & A.S. Rodrigues	x	-	MN652876
BHO_153	<i>B. ruderatus</i>	F	1	Spain	Madrid, Sierra de Guadarrama, Miraflores de La Sierra	40.800278	-3.756667	26/05/2014	V.L. Nunes, T.G. Laurentino & E. Marabuto	x	-	MN652877

s and GenBank accession numbers); outputs of RADseq analyses (number of initial and

RAD analysis	RAD sequencing lane	Barcode	Total reads	Retained reads (Process_Radtags)	Mapped reads properly paired (Bowtie2)	% of mapped reads	Number of loci (Stacks)	Mean Coverage (Stacks)	Samples of dataset BT_OUT	Samples of dataset BT_SSP	Samples of dataset BT_BTL	Frequency of missing data, dataset BT_OUT (VCFTools)	Frequency of missing data, dataset BT_SSP (VCFTools)	Frequency of missing data, dataset BT_BTL (VCFTools)
x	Lane 2	TTAAT	8109844	7734007	4605896	59.6	207938	17.92	x	x	x	0.00545171	0.0243446	0.0313586
removed	Lane 2	TGGTT	236766	120102	10810	9.0	747	8.77						
x	Lane 2	ATTAG	7315526	7127626	4316556	60.6	200433	17.62	x	x	x	0.0103193	0.0309482	0.0373926
x	Lane 2	AGGAC	4660438	4538676	2643980	58.3	136442	15.69	x	x	x	0.0564642	0.10408	0.105595
x	Lane 2	AGAGT	11736928	11350762	6998274	61.7	282072	19.6	x	x	x	0.00564642	0.0222748	0.0210276
x	Lane 2	AAGGG	5382908	5200161	3092488	59.5	148671	16.53	x	x	x	0.036215	0.0824956	0.0830133

x	Lane 1	CCGGT	7346304	7111056	4258414	59.9	150611	19.65	x	x	x	0.00778816	0.0283856	0.0291644
x	Lane 1	CAGTC	8489710	8257735	5231910	63.4	159071	21.98	x	x	x	0.00584112	0.0172482	0.0149936
removed	Lane 1	GTCAC	1968792	1872510	1055220	56.4	57562	12.11						
x	Lane 1	CTCTT	9251988	8770867	4913908	56.0	166760	20.57	x	x	x	0.00428349	0.0214863	0.0206619
x	Lane 1	TCTCT	15162584	14508026	8652690	59.6	208682	27.55	x	x	x	0.00175234	0.00818056	0.00895959
x	Lane 1	TATAC	3328796	3176285	1883466	59.3	74807	15.79	x	x	x	0.0630841	0.111275	0.109435
x	Lane 1	TGTGG	16898876	16338434	9724544	59.5	233263	27.95	x	x	x	0.00097352	0.00857481	0.00895959
x	Lane 1	CAACT	10468942	10070127	6041328	60.0	197828	21.49	x	x	x	0.00506231	0.015277	0.0152679
x	Lane 1	TTTTA	14787532	14167207	8373646	59.1	202131	27.48	x	x	x	0.00311526	0.008082	0.00722253
x	Lane 2	AAAAA	29499926	28594514	17801252	62.3	624304	22.68	x	x	x	0.00194704	0.00965898	0.00950814
removed	Lane 2	TCAGA	1399906	1244947	636054	51.1	41413	11.61						
x	Lane 2	TCGAG	4576352	4431323	2658398	60.0	134666	15.82	x	x	x	0.0609424	0.105855	0.109618
x	Lane 2	TGACC	7852030	7596370	4624002	60.9	211407	17.89	x	x	x	0.0132399	0.0361719	0.0326385
removed	Lane 2	ACGTA	337128	286528	152578	53.3	11331	9.29						
x	Lane 2	AACCC	13519934	13186619	8314690	63.1	325446	20.19	x	x	x	0.00097352	0.00798344	0.00987383

x	Lane 2	AATT	3466684	3201722	1756018	54.8	100208	14	x	x	x	0.138629	0.214567	0.224538
x	Lane 2	CAGTC	7963274	7719129	4557522	59.0	210993	17.74	x	x	x	0.0109034	0.0351863	0.0339184
x	Lane 2	CAACT	4352780	3997355	2198672	55.0	117516	14.95	x	x	x	0.0858645	0.149024	0.15149
x	Lane 2	CCAAC	8593044	8391856	5424226	64.6	229334	18.62	x	x	x	0.0107087	0.0249359	0.0232218
x	Lane 2	ACCAT	12321256	11796388	7165928	60.7	300695	19.57	x	x	x	0.00311526	0.0156712	0.0136222
x	Lane 2	ACTGC	10753424	10501149	6708582	63.9	277030	19.33	x	x	x	0.00584112	0.0189237	0.0165478
x	Lane 2	TCCG	7266592	7040446	4228904	60.1	195504	17.63	x	x	x	0.024338	0.0508575	0.0457122
x	Lane 2	TGTGG	9003316	8749325	5598210	64.0	239485	18.62	x	x	x	0.0122664	0.024246	0.0235875
x	Lane 2	TTTTA	3092968	2861849	1641620	57.4	92469	13.84	x	x	x	0.163162	0.237138	0.244012
removed	Lane 2	TGCAA	531410	428555	200060	46.7	13988	10.18						

x	Lane 2	GGCCT	6162856	5893962	3516654	59.7	170923	16.7	x	x	x	0.0272586	0.0613049	0.0571402
removed	Lane 2	GCTAA	1227456	1127861	658826	58.4	44981	10.83						
x	Lane 2	GCCGG	4709680	4580094	2737926	59.8	138878	15.87	x	x	x	0.0521807	0.0958999	0.0989212
x	Lane 2	TCCTC	12811682	12436736	7682970	61.8	302233	20.02	x	x	x	0.0155763	0.0408043	0.0417809
x	Lane 2	TATAC	2652990	2545204	1452420	57.1	85674	13.38	x	x	x	0.196262	0.266805	0.282319
x	Lane 2	TCTCT	5151776	4804789	2861534	59.6	144522	15.92	x	x	x	0.0488707	0.0885078	0.0911501

RAD analysis	RAD sequencing lane	Barcode	Total reads	Retained reads (Process_Radtags)	Mapped reads properly paired (Bowtie2)	% of mapped reads	Number of loci (Stacks)	Mean Coverage per individual (Stacks)	Samples of dataset BT_OUT	Samples of dataset BT_SSP	Samples of dataset BT_BTL	Frequency of missing data, dataset BT_OUT (VCFTools)	Frequency of missing data, dataset BT_SSP (VCFTools)	Frequency of missing data, dataset BT_BTL (VCFTools)
x	Lane 2	AGCTG	6885900	6683291	4088484	61.2	188241	17.73	x	x		0.0132399	0.0342007	
x	Lane 1	TTCCG	6060962	5863768	3366584	57.4	117490	18.99	x	x		0.00584112	0.0308496	
x	Lane 1	CCCCA	7457604	7241194	4306960	59.5	136690	20.94	x	x		0.00837227	0.0230633	
x	Lane 2	AGTCA	14513666	14223980	7884990	55.4	313539	20.05	x	x		0.00292056	0.0150798	
x	Lane 2	GACTA	7370284	7201198	4162876	57.8	198189	17.32	x	x		0.0144081	0.0381431	

RAD analysis	RAD sequencing lane	Barcode	Total reads	Retained reads (Process_Radtags)	Mapped reads properly paired (Bowtie2)	% of mapped reads	Number of loci (Stacks)	Mean Coverage per individual (Stacks)	Samples of dataset BT_OUT	Samples of dataset BT_SSP	Samples of dataset BT_BTL	Frequency of missing data, dataset BT_OUT (VCFTools)	Frequency of missing data, dataset BT_SSP (VCFTools)	Frequency of missing data, dataset BT_BTL (VCFTools)
x	Lane 2	CCCCA	6654410	6373456	1857064	29.1	94848	16.91	x			0.03602		
x	Lane 1	TCCTC	9630278	9237789	3797870	41.1	129379	21.73	x			0.122079		
x	Lane 2	CTCTT	7673874	7385489	3059096	41.4	138893	18.26	x			0.127531		
x	Lane 2	CTTCC	9673764	9393983	3976006	42.3	177966	19.07	x			0.120911		
x	Lane 1	GGCCT	5369388	4996943	2764028	55.3	100492	17.97	x			0.0216121		
removed	Lane 2	GTCAC	623266	527856	263090	49.8	18635	9.97						
removed	Lane 1	TGCAA	1606326	1471245	821262	55.8	44092	12.09						
removed	Lane 1	TACGT	2535204	2160906	743172	34.4	39070	11.54						
x	Lane 2	CGTAT	6366508	4132680	1413486	34.2	77692	15.43	x			0.178349		
x	Lane 2	CGCGC	6155140	5978181	2104318	35.2	107737	16.99	x			0.132788		
x	Lane 2	CCTTG	12943046	12573453	4313748	34.3	181112	20.31	x			0.118769		

x	Lane 2	CACAG	22264514	21811529	7183830	32.9	275076	22.88	x			0.0149922		
x	Lane 2	CATGA	7015302	6827666	2053094	30.1	107354	17.18	x			0.0428349		

Table S1b. List of samples with sequences downloaded from GenBank, with information concerning: sampling (taxa, sex, country, number of samples, origin, date of collection and bibliographic reference); outputs of mitochondrial COI analyses (haplotypes and GenBank accession numbers).

Taxa	Sex	n	Country / Island	Sampling Location	Latitude	Longitude	Date of collection	Collector(s)	mtDNA analysis	COI haplotype	GenBank accession no.
<i>B. terrestris</i>	F	5	Ireland	Wexford	52°20020 N	6°27036 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	5	Ireland	Cork	51°53052 N	8°28015 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	4	Ireland	Donegal	54°39016 N	8°06038 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	1	Ireland	Donegal	54°39016 N	8°06038 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	2	Ireland	Galway	53°16025 N	9°03006 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	3	Ireland	Galway	53°16025 N	9°03006 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	2	Ireland	Meath	53°37052 N	6°47027 W	01/06/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	3	Ireland	Meath	53°37052 N	6°47027 W	01/06/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Ireland	Kildare	53°09035 N	6°54032 W	01/06/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	5	Great Britain	North Wales, Rhyl	53°19010 N	3°29019 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	3	Great Britain	South Wales, Lampeter	51°18052 N	3°10049 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	2	Great Britain	South Wales, Lampeter	51°18052 N	3°10049 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	2	Isle of Man	Isle of Man	54°14009 N	4°32053 W	01/08/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	1	Isle of Man	Isle of Man	54°14009 N	4°32053 W	01/08/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Great Britain	England, Oxford	51°45008 N	1°15020 W	01/07/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	4	Great Britain	England, Dover	51°07046 N	1°18040 W	01/08/2007	Moreira <i>et al.</i> (2015)	x	HA / H8	KP670306
<i>B. terrestris</i>	F	1	Great Britain	England, Dover	51°07046 N	1°18040 W	01/08/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Greece	Macedonia	40°45000 N	22°53059 E	01/07/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Portugal	Bragança	41°48025 N	6°45033 W	01/05/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Croatia	Pula	44°52010 N	13°50028 E	01/03/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Croatia	Zadar	44°01046 N	17°48000 E	01/03/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	France	Samer	44°38021 N	1°44042 E	01/07/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	France	Beauvais	49°25045 N	2°04051 E	01/07/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	France	Grenoble	45°11039 N	5°43053 E	01/05/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Poland	Grudziadz	54°20038 N	18°39056 E	01/06/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Poland	Katowice	50°15053 N	19°01025 E	01/06/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Denmark	Bilund	56°15050 N	9°30006 E	01/08/2007	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. terrestris</i>	F	5	Latvia	Jelgava	56°56047 N	24°06017 E	07/05/2008	Moreira <i>et al.</i> (2015)	x	HB / H1	KP670307
<i>B. t. sassaricus</i>	M	7	Sardinia	--	40°47'27"N	08°03'12"E	--	Coppée (2010)	x	H11	JQ769086
<i>B. t. terrestris</i>	M	4	France (Southeast)	--	43°25'32"N	06°25'23"E	--	Coppée (2010)	x	H1	JQ769069
<i>B. t. terrestris</i>	M	4	France (North)	--	48°50'37"N	02°21'35"E	--	Coppée (2010)	x	H1	JQ769080
<i>B. t. terrestris</i>	M	4	France (Southwest)	--	42°28'38"N	01°55'04"E	--	Coppée (2010)	x	H1	JQ769088
<i>B. t. terrestris</i>	M	4	Belgium	--	50°50'22"N	04°23'44"E	--	Coppée (2010)	x	H1	JQ769070
<i>B. t. terrestris</i>	M	3	Germany	--	51°56'27"N	07°33'03"E	--	Coppée (2010)	x	H13	JQ769081
<i>B. t. terrestris</i>	M	3	Denmark	--	54°59'01"N	12°00'32"E	--	Coppée (2010)	x	H1	JQ769082
<i>B. t. terrestris</i>	M	5	Sweden	--	59°51'43"N	17°38'00"E	--	Coppée (2010)	x	H1	JQ769083
<i>B. t. xanthopus</i>	M	5	Corsica	--	42°17'25"N	08°52'40"E	--	Coppée (2010)	x	H15	JQ769071
<i>B. t. dalmatinus</i>	M	5	Greece (Crete)	--	35°23'28"N	23°34'30"E	--	Coppée (2010)	x	H1	JQ769072
<i>B. t. dalmatinus</i>	M	4	Turkey	--	39°55'54"N	32°51'48"E	--	Coppée (2010)	x	H1	JQ769078
<i>B. t. dalmatinus</i>	M	4	France (Southeast)	--	43°25'32"N	06°25'23"E	--	Coppée (2010)	x	H1	JQ769079
<i>B. t. lusitanicus</i>	M	3	Madeira	--	32°48'01"N	16°50'43"W	--	Coppée (2010)	x	H1	JQ769073
<i>B. t. lusitanicus</i>	M	3	France (Southwest)	--	42°28'51"N	01°55'25"E	--	Coppée (2010)	x	H1	JQ769087
<i>B. t. lusitanicus</i>	M	5	Spain	--	36°42'59"N	04°25'03"W	--	Coppée (2010)	x	H1	JQ769076

<i>B. t. africanus</i>	M	5	Morocco	--	35°03'40"N	05°09'60"W	--	Coppée (2010)	x	H16	JQ769074
<i>B. t. calabricus</i>	M	5	Italy	--	35°59'35"N	13°52'49"E	--	Coppée (2010)	x	H1	JQ769075
<i>B. t. audax</i>	M	5	Ireland	--	52°57'03,3"N	9°04'41,8"W	--	Coppée (2010)	x	H2	JQ769077
<i>B. t. audax</i>	M	3	Great Britain (North)	--	51°05'43,0"N	0°38'33,2"E	--	Coppée (2010)	x	H8	JQ769084
<i>B. t. audax</i>	M	5	Great Britain (Southwest)	--	50°21'33"N	03°50'15"W	--	Coppée (2010)	x	H8	JQ769085
<i>B. terrestris</i>	M	1	Great Britain	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H8	JQ843647
<i>B. terrestris</i>	M	1	Great Britain	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H8	JQ843646
<i>B. terrestris</i>	F	1	Turkey	Sinop	--	--	2010	Williams <i>et al.</i> (2012b)	x	H1	JQ843645
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843658
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843669
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H12	JQ843663
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843643
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843664
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843644
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843611
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843612
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843613
<i>B. terrestris</i>	F	1	Turkey	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843614
<i>B. terrestris</i>	F	1	Algeria	Bouira	--	--	2005	Williams <i>et al.</i> (2012b)	x	H14	JQ843667
<i>B. terrestris</i>	F	1	Italy	Siena	--	--	1995	Williams <i>et al.</i> (2012b)	x	H14	JQ843662
<i>B. terrestris</i>	F	1	Spain	Canary Islands	--	--	2008	Williams <i>et al.</i> (2012b)	x	H17	JQ843661
<i>B. terrestris</i>	F	1	Spain	Canary Islands	--	--	2008	Williams <i>et al.</i> (2012b)	x	H17	JQ843660
<i>B. terrestris</i>	F	1	Spain	Canary Islands	--	--	2008	Williams <i>et al.</i> (2012b)	x	H17	JQ843659
<i>B. terrestris</i>	F	1	France	Perpignan	--	--	2008	Williams <i>et al.</i> (2012b)	x	H1	JQ843670
<i>B. terrestris</i>	F	1	Latvia	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843609
<i>B. terrestris</i>	F	1	Latvia	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843610
<i>B. terrestris</i>	F	1	Great Britain	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H8	JQ843628
<i>B. terrestris</i>	F	1	Great Britain	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H8	JQ843631
<i>B. terrestris</i>	F	1	Great Britain	Bromley, close to London	--	--	2011	Williams <i>et al.</i> (2012b)	x	H8	JQ843632
<i>B. terrestris</i>	F	1	Germany	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843633
<i>B. terrestris</i>	F	1	Great Britain	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H8	JQ843634
<i>B. terrestris</i>	F	1	Greece	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843635
<i>B. terrestris</i>	F	1	Sweden	Uppsala	--	--	2011	Williams <i>et al.</i> (2012a)	x	H1	JQ692970
<i>B. terrestris</i>	F	1	Sweden	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843636
<i>B. terrestris</i>	F	1	Sweden	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843637
<i>B. terrestris</i>	F	1	Sweden	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843638
<i>B. terrestris</i>	F	1	Switzerland	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843639
<i>B. terrestris</i>	F	1	Switzerland	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843640
<i>B. terrestris</i>	F	1	Switzerland	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843641
<i>B. terrestris</i>	F	1	Switzerland	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H12	JQ843642
<i>B. terrestris</i>	F	1	Switzerland	--	--	--	--	Williams <i>et al.</i> (2012b)	x	H1	JQ843629
<i>B. terrestris</i>	--	1	Germany	Brandenburg, Mallnow	52.56 N	14.48E	28/03/2012	Schmidt <i>et al.</i> (2015)	x	H12	KJ838396
<i>B. terrestris</i>	--	1	Germany	Brandenburg, Mallnow	52.56 N	14.48E	28/03/2012	Schmidt <i>et al.</i> (2015)	x	H1	KJ838178
<i>B. terrestris</i>	--	1	Germany	Baden-Wuerttemberg, Muehl	47.82 N	7.62 E	07/04/2012	Schmidt <i>et al.</i> (2015)	x	H1	KJ838879

Table S2. The sample size and diversity indices across *B. t. lusitanicus* samples by Iberian region considering mitochondrial DNA COI data and RADseq dataset BT_BT_L. Note that six individuals were excluded from RADseq dataset. n , number of individuals analysed; h , average haplotype diversity; π , average nucleotide diversity; H_o , observed heterozygosity; H_E , expected heterozygosity.

Geographic region	COI			RADseq			
	n (total = 198)	h	π	n (total = 31)	H_o	H_E	π
IP-NW	24 (2 hap.)	0.08330	0.00014	2	--	--	--
IP-WE	56 (2 hap.)	0.07010	0.00012	11	0.2696	0.2919	0.2706
IP-SW	20 (2 hap.)	0.10000	0.00034	3	--	--	--
IP-NO	28 (2 hap.)	0.07140	0.00012	2	--	--	--
IP-CE	32 (1 hap.)	0	0	5	0.3148	0.3600	0.2702
IP-SE	38 (5 hap.)	0.19700	0.00044	8	0.2537	0.3128	0.2706

STACKS parameter testing

Methods

Preliminary tests of parameter values for each STACKS module were carried out in order to identify the optimal parameter values for use in the final STACKS analysis. The three datasets (BT_OUT, BT_SSP and BT_BTL) were tested separately. For the three first modules – *pstacks*, *cstacks* and *sstacks* – tests were performed using three sets of three randomly chosen individuals, each set corresponding to each one of the three datasets. For the last module – *populations* – all the samples were used because the number of SNPs obtained vary with population size and number.

Pstacks extract stacks that have been aligned to a reference genome and identify SNPs in each individual (Catchen *et al.*, 2013). In this module, the parameter -m (minimum depth of coverage to report a stack) was tested. Several tests were performed by increasing the ‘m’ value in increments of 2, from 2 to 10, while non-test parameters were kept as default. For each test, the number of created loci and respective mean coverage was evaluated. The input for *cstacks* module was created by running *pstacks* with the selected “optimal” parameters. *Cstacks* merges loci from multiple individuals, creating a set of consensus loci or catalog (Catchen *et al.*, 2013). For the module, the parameter -n (maximum number of mismatches allowed between homologous loci among individuals when building the catalog) was also tested by increasing the value in increments of 2 from 2 to 10, while keeping non-test parameters as default. The number of loci added to the catalog for each ‘n’ value, as well as the number of verified matches against the catalog was then evaluated.

Populations is used to compute population-level summary statistics and to output SNPs according to some parameters (Catchen *et al.*, 2013). In this module two main parameters were tested: -p, which represents the minimum number of populations that a locus must be present in for it to be retained in the final dataset; -r, which represents the minimum percentage of individuals in a population that a locus must be present in for it to be retained in the final dataset (Catchen *et al.*, 2013). Several combinations using these two parameters were tested as follows: ‘p’ was set at values between 3 and 6 for dataset BT_BTL, at values between 5 and 9 for dataset BT_SSP and at values between 7 and 14 for dataset BT_OUT, which corresponds to 50% of taxa or above, for each dataset; ‘r’ was set to 50% and 100% for each ‘p’ setting. Only these two values of ‘r’ were tested because some taxa are only represented by two or less individuals. After testing each combination of parameters (-r; -p), we evaluated the number of retained SNP loci in each final dataset, the average coverage per SNP per sample and the average frequency of missing data per sample, using VCFTOOLS version 0.1.14.

Results

Figure S1 shows the results of parameter tests for *pstacks*. In *pstacks*, an ‘m’ parameter value of two (minimum of two reads to report a stack) resulted in a very large number of loci (min: 405 555, máx: 1 700 054), and with a low coverage, as expected (min: 7.26, máx 10.57). Increasing the minimum stack depth parameter to four prevented reads with possible sequencing errors from forming stacks and the number of

loci decreases considerably, as mean coverage increases. After $m=6$ (number of loci: min: 117 490, máx: 624 304; mean coverage: min: 17.73, máx: 27.95), further increases in 'm' resulted in slowed changes in number of loci, while coverage increases considerably, probably because the rate at which stacks absent from the reference were removed slowed. After this value, true stacks began to be dismantled, resulting in a lower number of loci, while the stacks that are maintained have very high coverage (Catchen *et al.*, 2013). The results from $m=6$ seem to be the ones which result in the best combination between loci number and coverage. In this way, the value of m set to 6 was chosen to perform the final STACKS analysis. The output results obtained from the parameter test using $m=6$ were used as input for the *cstacks* parameter tests.

Figure S2 shows the results of parameter tests for *cstacks* and *sstacks*. Increments in the maximum number of mismatches allowed between homologous loci from multiple individuals when building the catalog, seem to have a very small impact in the number of loci added to the catalog and in the number of verified matches of an individual against the catalog. This value varies with the evolutionary distance of the members being examined, and the small variation observed is probably the result of the small divergence among individuals (Catchen *et al.*, 2011). The value $n=2$ was chosen for final STACKS analysis and for running the input for *populations* parameters tests.

Figure S3 shows the results of parameter tests for *populations* module. Results for $r=50\%$ typically showed higher numbers of SNP loci with lower coverage but also with lower amount of missing data. This means that few SNPs are present in all individuals of each population. Higher values of 'p' typically yielded lower numbers of loci but with higher coverage and also lower amount of missing data. This means that, as the stringency of 'p' increased, less SNPs are available, but they have a relatively good coverage across all samples, and as a consequence, less missing data. Among the different tests, the number of retained loci ranged from 40 045 (average coverage per SNP per sample: 50.70; average frequency of missing data per sample: 0.10) to 14 231 for dataset BT_BTL (average coverage per SNP per sample: 73.25; average frequency of missing data per sample: 0), from 41 849 (average coverage per SNP per sample: 49.7; average frequency of missing data per sample: 0.1) to 14 580 (average coverage per SNP per sample: 72.6; average frequency of missing data per sample: 0) for dataset BT_SSP and from 45 184 (average coverage per SNP per sample: 44.69; average frequency of missing data per sample: 0.20) to 3 772 (average coverage per SNP per sample: 75.18; average frequency of missing data per sample: 0) for dataset BT_OUT. Based on these results, outputs from runs using the parameters p5r50 (dataset BT_BTL), p8r50 (dataset BT_SSP) and p13r50 (dataset BT_OUT) retain the better combination of number of loci, average coverage per SNP per sample and amount of missing data and these parameters were chosen for final STACKS analysis.




Figure S1. Results of parameter tests for the STACKS module *pstacks*. (a) Changes in number of created loci (*pstacks*) and (b) loci mean coverage obtained (*pstacks*) for each value of ‘*m*’ tested for three randomly selected individuals from each dataset.




Figure S2. Results of parameter tests for the STACKS modules *ctsacks* and *sstacks*. (a) Changes in number of loci added to the catalog (*ctsacks*) and (b) number of matches against the catalog (*sstacks*) for each value of ‘*n*’ tested for three randomly selected individuals from each dataset.




Figure S3. Results of parameter tests for the STACKS module *populations*. Variation in number of SNP loci in the final dataset, in average coverage per SNP and per sample and in average frequency of missing data per sample obtained for each combination of parameters (-r; -p) tested in *populations* module, using (a) dataset BT_BTL, (b) dataset BT_SSP and (c) dataset BT_OUT.


```
#####
# List of command line commands used for RADseq dataset analyses ##
#####
# in the commands below
# input_files_path = input file path
# output_files_path = output path to write results
# output_directory_path = output path to directory where to write the results
# barcodes_file_path = path to a file containing barcodes
# log_files_path = path to write logfile
# list_of_samples = list of the files corresponding to each sample
# populations_file_path = path to a file containing the list of populations

### Build index using BOWTIE2 version 2.1.0
bowtie2-build -f input_files_path Bt_all

### Prepare reads (cleaning and demultiplexing) with STACKS version 1.29 - pipeline
process_radtags
process_radtags -1 input_files_path -2 input_files_path -o output_files_path -b
barcodes_file_path -c -q -r -e pstI -E phred33 -i gzfastq -D

### Convert barcodes to sample name
#!/bin/bash
mv input_files_path/sample_ATTAG.1.fq.gz output_files_path/BTL_236.1.fq.gz
...

### Align reads with reference genome using BOWTIE2 version 2.1.0 #
#!/bin/bash

samples="
list_of_samples
"

for sample in $samples
do

bowtie2 --trim3 3 --sensitive -p 6 -x input_files_path/Bt_all -1 input_files_path/$
{sample}.1.fq.gz -2 input_files_path/${sample}.2.fq.gz -S input_files_path/$
{sample}_Bt_all.sam

done
```

```
### Conver SAM files to BAM using SAMTOOLS version 0.1.19
```

```
#!/bin/bash
```

```
for samfile in $(ls input_files_path/*.sam); do samtools view -Sb ${samfile} > ${samfile}.bam; done
```

```
### Sort and filter BAM files using SAMTOOLS version 0.1.19
```

```
#!/bin/bash
```

```
samples=""
```

```
list_of_samples
```

```
"
```

```
for sample in $samples
```

```
do
```

```
samtools sort input_files_path/${sample}_Bt_all.sam.bam output_files_path/${sample}_Bt_all.sam.bam.sorted
```

```
samtools index input_files_path/${sample}_Bt_all.sam.bam.sorted.bam
```

```
samtools view -q 20 -F 0x0004 -b input_files_path/${sample}_Bt_all.sam.bam.sorted.bam > output_files_path/${sample}_Bt_all.sorted.quality.bam
```

```
samtools flagstat input_files_path/${sample}_Bt_all.sorted.quality.bam > output_files_path/${sample}_Bt_all.sorted.quality_flagstat
```

```
done
```

```
### Build loci from ref using STACKS version 1.45 - pipeline pstacks
```

```
#!/bin/bash
```

```
samples=""
```

```
list_of_samples
```

```
"
```

```
i=1
```

```
for sample in $samples
```

```
do
```

```
    pstacks -t bam -i $i -m 6 -p 4 \
```

```
        -f input_files_path/$sample \
```

```
        -o output_files_path &>> log_files_path/Log_pstacks
```

```
    let "i+=1";
```

```
done
```

```
### Create catalog and match samples to catalog using STACKS version 1.45 -  
pipeline cstacks and sstacks  
#!/bin/bash
```

```
samples="  
list_of_samples  
"
```

```
samp=""  
for sample in $samples  
do  
    samp+="-s input_files_path/$sample ";  
done
```

```
cstacks -p 4 -b 1 -n 2 -o output_files_path $samp &>> log_files_path/Log_cstacks
```

```
for sample in $samples  
do  
    sstacks -p 4 -b 1 -c input_files_path/batch_1 \  
    -s input_files_path/${sample} \  
    -o output_files_path &>> log_files_path/Log_sstacks  
done
```

```
### Filter and export SNPs to VCF file format using STACKS version 1.45 - pipeline  
populations, using dataset BT_OUT  
#!/bin/bash
```

```
populations -b 1 -P input_files_path -M input_files_path -t 4 -p 13 -r 50 --vcf --  
write_random_snp
```

```
### Filter and export SNPs to VCF file format using STACKS version 1.45 - pipeline  
populations, using dataset BT_SSP  
#!/bin/bash
```

```
populations -b 1 -P input_files_path -M input_files_path -t 4 -p 8 -r 50 --vcf --  
write_random_snp
```

```
### Filter and export SNPs to VCF file format using STACKS version 1.45 - pipeline  
populations, using dataset BT_BTL  
#!/bin/bash
```

```
populations -b 1 -P input_files_path -M input_files_path -t 4 -p 5 -r 50 --vcf --  
write_random_snp
```

```
### Determine missing data per individual and filter VCF files using VCFTOOLS  
version 0.1.15
```

```
vcftools --vcf input_file_path --missing-indv  
vcftools --vcf input_file_path --max-missing 0.75 --maf 0.05 --recode --out  
output_file_path
```

```
### Determine per-SNP nucleotide diversity, observed and expected heterozygosities
```

```
vcftools --vcf input_file_path --site-pi  
vcftools --vcf input_file_path --het
```

```
### Principal component analysis of the three datasets using the R script  
snp_pca_static.R (https://github.com/CoBiG2/RAD\_Tools) as of commit 134b11d
```

```
snp_pca_static.R input_files_path output_files_path populations_file_path
```

```
### Run BAYESCAN version 2.1 using both datasets BT_SSP and BT_BTL  
./bayescan_2.1 input_file_path -snp -threads 14 -od output_directory_path
```

```
### Run SELESTIM version 1.1.7 using both datasets BT_SSP and BT_BTL  
./selestim -threads 12 -file input_files_path -outputs output_directory path -thin  
20 -npilot 50 -lpilot 1000 -burnin 100000 -length 1000000 -calibration
```

```
### Run Maverick version 1.0.4 with Structure_threader version 1.2.4, using dataset  
BT_SSP
```

```
~/local/bin/structure_threader run -K 5 -i input_file_path -o output_file_path -t  
10 -mv ~/local/bin/Maverick --params parameters.txt --log=1
```

```
### Run Maverick version 1.0.4 with Structure_threader version 1.2.4, using dataset  
BT_BTL
```

```
~/local/bin/structure_threader run -K 4 -i input_file_path -o output_file_path -t  
10 -mv ~/local/bin/Maverick --params parameters.txt --log=1
```

Final set of parameters (parameters.txt) used for run Maverick, using dataset BT_SSP

Data properties

headerRow_on t
popCol_on f
ploidyCol_on f
ploidy 2
missingData -9

Model parameters

Kmin 1
Kmax 5
admix_on t
fixAlpha_on f
alpha 1.0,0.845,0.448,0.414,0.32
alphaPropSD 1.0,0.2125,0.0975,0.065,0.063

Simulation parameters

exhaustive_on f
mainRepeats 10
mainBurnin 5000
mainSamples 50000
thermodynamic_on t
thermodynamicRungs 50
thermodynamicBurnin 5000
thermodynamicSamples 50000
EMalgorithm_on f
EMrepeats 100
EMiterations 100

Basic output properties

outputLog_on t
outputLikelihood_on t
outputQmatrix_ind_on t
outputQmatrix_pop_on f
outputQmatrixError_ind_on t
outputQmatrixError_pop_on f
outputEvidence_on t
outputEvidenceNormalised_on t
outputEvidenceDetails_on t

Final set of parameters (parameters.txt) used for run Maverick, using dataset BT_BTL

Data properties

headerRow_on t
popCol_on f
ploidyCol_on f

```
# ploidy      2
# missingData -9
#### Model parameters
# Kmin 1
# Kmax 4
# admix_on    t
# fixAlpha_on f
# alpha      1.0, 3.94, 1.58, 1.05
# alphaPropSD 1.0, 0.51, 0.18, 0.12
#### Simulation parameters
# exhaustive_on    f
# mainRepeats 10
# mainBurnin 5000
# mainSamples 50000
# thermodynamic_on t
# thermodynamicRungs      50
# thermodynamicBurnin     5000
# thermodynamicSamples     50000
# EMalgorithm_on    f
# EMrepeats  100
# EMiterations      100
#### Basic output properties
# outputLog_on      t
# outputLikelihood_on      t
# outputQmatrix_ind_on      t
# outputQmatrix_pop_on      f
# outputQmatrixError_ind_on      t
# outputQmatrixError_pop_on      f
# outputEvidence_on t
# outputEvidenceNormalised_on      t
# outputEvidenceDetails_on t
```